


**FUSED HETEROARYL DERIVATIVES FOR USE AS P38 KINASE
INHIBITORS IN THE TREATMENT OF I.A. RHEUMATOID ARTHRITIS**

Rec'd PCT/PTC 31 JAN 2005

This invention relates to novel compounds and their use as pharmaceuticals, particularly as p38 kinase inhibitors, for the treatment of conditions or disease states mediated by p38 kinase activity or mediated by cytokines produced by the activity of p38 kinase.

We have now found a group of novel compounds that are inhibitors of p38 kinase.

According to the invention there is provided a compound of formula (I):



(I)

wherein

A is a fused 5-membered heteroaryl ring optionally substituted by up to two substituents independently selected from C₁₋₆alkyl, -(CH₂)_m-C₃₋₇cycloalkyl, halogen, cyano, trifluoromethyl, -(CH₂)_mOR³, -(CH₂)_mCO₂R³, -(CH₂)_mNR³R⁴, -(CH₂)_mCONR³R⁴, -(CH₂)_mNHCOR³, -(CH₂)_mSO₂NR³R⁴, -(CH₂)_mNHSO₂R³, -(CH₂)_mSO₂(CH₂)_nR⁵, a 5- or 6-membered heterocyclyl ring containing nitrogen optionally substituted by C₁₋₂alkyl or -(CH₂)_mCO₂R³, and a 5-membered heteroaryl ring optionally substituted by C₁₋₂alkyl;

R¹ is selected from methyl and chloro;

R² is selected from -NH-CO-R⁶ and -CO-NH-(CH₂)_q-R⁷;

R³ is selected from hydrogen, C₁₋₆alkyl optionally substituted by up to two OH groups, -(CH₂)_m-C₃₋₇cycloalkyl, -(CH₂)_mphenyl optionally substituted by R⁸ and/or R⁹ and -(CH₂)_mheteroaryl optionally substituted by R⁸ and/or R⁹,

R⁴ is selected from hydrogen and C₁₋₆alkyl, or

R³ and R⁴, together with the nitrogen atom to which they are bound, form a 5- or 6-membered heterocyclic ring optionally containing one additional heteroatom selected from oxygen, sulfur and N-R¹⁰;

R⁵ is selected from C₁₋₆alkyl optionally substituted by up to three halogen atoms, C₂₋₆alkenyl optionally substituted by phenyl, C₃₋₇cycloalkyl, heteroaryl optionally substituted by up to three R⁸ and/or R⁹ groups, and phenyl optionally substituted by R⁸ and/or R⁹;

R⁶ is selected from hydrogen, C₁₋₆alkyl, -(CH₂)_q-C₃₋₇cycloalkyl, trifluoromethyl, -(CH₂)_rheteroaryl optionally substituted by R¹¹ and/or R¹², and -(CH₂)_rphenyl optionally substituted by R¹¹ and/or R¹²;

R⁷ is selected from hydrogen, C₁₋₆alkyl, C₃₋₇cycloalkyl, -CONHR¹³, phenyl optionally substituted by R¹¹ and/or R¹², and heteroaryl optionally substituted by R¹¹ and/or R¹²;

R⁸ and R⁹ are each independently selected from halogen, cyano, trifluoromethyl, nitro, C₁₋₆alkyl, C₁₋₆alkoxy, -CONR¹³R¹⁴, -COR¹⁵, -CO₂R¹⁵, and heteroaryl, or

R⁸ and R⁹ are linked to form a fused 5-membered heterocyclyl ring containing one heteroatom selected from oxygen, sulphur and N-R¹⁰, or a fused heteroaryl ring;

R¹⁰ is selected from hydrogen and methyl;

R¹¹ is selected from C₁₋₆alkyl, C₁₋₆alkoxy, -(CH₂)_q-C₃₋₇cycloalkyl, -CONR¹³R¹⁴, -NHCOR¹⁴, halogen, CN, -(CH₂)_sNR¹⁶R¹⁷, trifluoromethyl, phenyl optionally substituted by one or more R¹² groups, and heteroaryl optionally substituted by one or more R¹² groups;

R¹² is selected from C₁₋₆alkyl, C₁₋₆alkoxy, halogen, trifluoromethyl, and -(CH₂)_sNR¹⁶R¹⁷;

R¹³ and R¹⁴ are each independently selected from hydrogen and C₁₋₆alkyl, or R¹³ and R¹⁴, together with the nitrogen atom to which they are bound, form a 5- or 6-membered heterocyclic ring optionally containing one additional heteroatom selected from oxygen, sulfur and N-R¹⁰, wherein the ring may be substituted by up to two C₁₋₆alkyl groups;

R¹⁵ is C₁₋₆alkyl;

R¹⁶ is selected from hydrogen, C₁₋₆alkyl and -(CH₂)_q-C₃₋₇cycloalkyl optionally substituted by C₁₋₆alkyl,

R¹⁷ is selected from hydrogen and C₁₋₆alkyl, or

R¹⁶ and R¹⁷, together with the nitrogen atom to which they are bound, form a 5- or 6-membered heterocyclic ring optionally containing one additional heteroatom selected from oxygen, sulfur and N-R¹⁰;

X and Y are each independently selected from hydrogen, methyl and halogen;

m is selected from 0, 1, 2 and 3;


n is selected from 0, 1, 2 and 3;

q is selected from 0, 1 and 2;

r is selected from 0 and 1; and

s is selected from 0, 1, 2 and 3.

According to a further embodiment of the invention there is provided a compound of formula (IA):



(IA)

wherein

- A is a fused 5-membered heteroaryl ring optionally substituted by up to two substituents independently selected from C₁₋₆alkyl, -(CH₂)_m-C₃₋₇cycloalkyl, halogen, cyano, trifluoromethyl, -(CH₂)_mOR³, -(CH₂)_mNR³R⁴, -(CH₂)_mCONR³R⁴, -(CH₂)_mNHCOR³, -(CH₂)_mSO₂NR³R⁴, -(CH₂)_mNHSO₂R³, -(CH₂)_mSO₂(CH₂)_nR⁵, a 5- or 6-membered heterocyclyl ring containing nitrogen optionally substituted by C₁₋₂alkyl and a 5-membered heteroaryl ring optionally substituted by C₁₋₂alkyl;
- R¹ is selected from methyl and chloro;
- R² is selected from -NH-CO-R⁶ and -CO-NH-(CH₂)_q-R⁷;
- R³ is selected from hydrogen, C₁₋₆alkyl optionally substituted by up to two OH groups, -(CH₂)_m-C₃₋₇cycloalkyl, -(CH₂)_mphenyl optionally substituted by R⁸ and/or R⁹ and -(CH₂)_mheteroaryl optionally substituted by R⁸ and/or R⁹
- R⁴ is selected from hydrogen and C₁₋₆alkyl, or
- R³ and R⁴, together with the nitrogen atom to which they are bound, form a 5- or 6-membered heterocyclic ring optionally containing one additional heteroatom selected from oxygen, sulfur and N-R¹⁰;
- R⁵ is selected from C₁₋₆alkyl, C₃₋₇cycloalkyl, heteroaryl optionally substituted by R⁸ and/or R⁹, and phenyl optionally substituted by R⁸ and/or R⁹;
- R⁶ is selected from hydrogen, C₁₋₆alkyl, -(CH₂)_q-C₃₋₇cycloalkyl, trifluoromethyl, -(CH₂)_rheteroaryl optionally substituted by R¹¹ and/or R¹², and -(CH₂)_rphenyl optionally substituted by R¹¹ and/or R¹²;
- R⁷ is selected from hydrogen, C₁₋₆alkyl, C₃₋₇cycloalkyl, CONHR¹³, phenyl optionally substituted by R¹¹ and/or R¹², and heteroaryl optionally substituted by R¹¹ and/or R¹²;
- R⁸ and R⁹ are each independently selected from halogen, cyano, trifluoromethyl, C₁₋₆alkyl, C₁₋₆alkoxy, COR¹⁵, CO₂R¹⁵, and heteroaryl, or
- R⁸ and R⁹ are linked to form a fused 5-membered heterocyclyl ring containing one heteroatom selected from oxygen, sulphur and N-R¹⁰;
- R¹⁰ is selected from hydrogen and methyl;
- R¹¹ is selected from C₁₋₆alkyl, C₁₋₆alkoxy, -(CH₂)_q-C₃₋₇cycloalkyl, -CONR¹³R¹⁴, -NHCOR¹⁴, halogen, CN, -(CH₂)_sNR¹⁶R¹⁷, trifluoromethyl, phenyl optionally substituted by one or more R¹² groups, and heteroaryl optionally substituted by one or more R¹² groups;

R^{12} is selected from C_{1-6} alkyl, C_{1-6} alkoxy, halogen, trifluoromethyl, and $-(CH_2)_sNR^{16}R^{17}$;

R^{13} and R^{14} are each independently selected from hydrogen and C_{1-6} alkyl, or

- 5 R^{13} and R^{14} , together with the nitrogen atom to which they are bound, form a 5- or 6-membered heterocyclic ring optionally containing one additional heteroatom selected from oxygen, sulfur and N- R^{10} , wherein the ring may be substituted by up to two C_{1-6} alkyl groups;

R^{15} is C_{1-6} alkyl;

- 10 R^{16} is selected from hydrogen, C_{1-6} alkyl and $-(CH_2)_q-C_{3-7}$ cycloalkyl optionally substituted by C_{1-6} alkyl,

R^{17} is selected from hydrogen and C_{1-6} alkyl, or

R^{16} and R^{17} , together with the nitrogen atom to which they are bound, form a 5- or 6-membered heterocyclic ring optionally containing one additional heteroatom selected from oxygen, sulfur and N- R^{10} ;

- 15 m is selected from 0, 1, 2 and 3;


n is selected from 0, 1, 2 and 3;

q is selected from 0, 1 and 2;

r is selected from 0 and 1; and


s is selected from 0, 1, 2 and 3.

- 20 Representative examples of A include fused 5-membered heteroaryl rings containing up to two heteroatoms independently selected from oxygen, nitrogen and sulfur. Preferred fused 5-membered heteroaryl rings include rings containing up to two heteroatoms independently selected from oxygen and nitrogen, in particular rings containing two heteroatoms selected from oxygen and nitrogen such as rings
- 25 containing a nitrogen atom and one additional heteroatom selected from oxygen and nitrogen. In one embodiment, A is selected from fused pyrazolyl and isoxazolyl rings such as those shown below:



30

In a further embodiment, A is selected from fused pyrrolyl and thiazolyl rings such as those shown below:



Ring A may be optionally substituted by up to two substituents, located on any position on the ring. Preferably ring A is substituted by one substituent.

- 5 In one embodiment, A is optionally substituted by up to two substituents independently selected from C₁₋₆alkyl, -(CH₂)_m-C₃₋₇cycloalkyl, halogen, cyano, trifluoromethyl, -(CH₂)_mOR³, -(CH₂)_mNR³R⁴, -(CH₂)_mCONR³R⁴, -(CH₂)_mNHCOR³, -(CH₂)_mSO₂NR³R⁴, -(CH₂)_mNHSO₂R³, -(CH₂)_mSO₂(CH₂)_nR⁵, a 5- or 6-membered heterocyclyl ring containing nitrogen
10 optionally substituted by C₁₋₂alkyl, and a 5-membered heteroaryl ring optionally substituted by C₁₋₂alkyl.

- Representative examples of substituents for A include C₁₋₄alkyl, in particular methyl; -(CH₂)_m-C₃₋₇cycloalkyl, in particular -(CH₂)_m-cyclopropyl; -(CH₂)_mCO₂R³; -(CH₂)_mNR³R⁴; -(CH₂)_mCONR³R⁴; -(CH₂)_mNHCOR³; -
15 (CH₂)_mSO₂(CH₂)_nR⁵; and a 5- or 6-membered heterocyclyl ring containing nitrogen, in particular 4-piperidinyl, optionally substituted by C₁₋₂alkyl or -(CH₂)_mCO₂R³.

- Typical substituents for A include C₁₋₄alkyl, in particular methyl, -(CH₂)_mOR³, -(CH₂)_mNR³R⁴, -(CH₂)_mSO₂(CH₂)_nR⁵, and a 5- or 6-membered
20 heterocyclyl ring containing nitrogen, in particular piperazinyl and piperidinyl such as 4-piperidinyl.

A representative example of R¹ is methyl.

In one embodiment, R² is -CO-NH-(CH₂)_q-R⁷.

- Representative examples of R³ include hydrogen; C₁₋₆alkyl optionally
25 substituted by up to two OH groups, in particular methyl, ethyl, n-propyl, isopropyl, t-butyl or 2,2-dimethylpropyl optionally substituted by up to two OH groups; -(CH₂)_m-C₃₋₇cycloalkyl, in particular -(CH₂)_m-cyclopropyl; -(CH₂)_mphenyl optionally substituted by R⁸ and/or R⁹; and -(CH₂)_mheteroaryl, in particular thiazolyl, optionally substituted by R⁸ and/or R⁹.

- 30 Further representative examples of R³ include hydrogen and C₁₋₄alkyl substituted by up to two OH groups, in particular 1,3-dihydroxyprop-2-yl.

Representative examples of R⁴ include hydrogen and C₁₋₄alkyl such as methyl. In particular, a representative example of R⁴ is hydrogen.

- In one embodiment, R³ and R⁴, together with the nitrogen atom to which they
35 are bound, form a 5- or 6-membered heterocyclic ring optionally containing one additional heteroatom selected from oxygen, sulfur and N-R¹⁰, in particular a pyrrolidinyl, piperidinyl, piperazinyl or 4-methylpiperazinyl, or morpholinyl ring.

In one embodiment, R⁵ is selected from C₁₋₆alkyl, C₃₋₇cycloalkyl, heteroaryl optionally substituted by R⁸ and/or R⁹, and phenyl optionally substituted by R⁸ and/or R⁹.

Representative examples of R⁵ include C₁₋₆alkyl optionally substituted by up to three halogen atoms, in particular methyl, ethyl, n-propyl, isopropyl, n-butyl and n-hexyl optionally substituted by up to three halogen atoms; C₂₋₆alkenyl optionally substituted by phenyl, in particular ethenyl optionally substituted by phenyl; C₃₋₇cycloalkyl, in particular cyclopropyl; heteroaryl optionally substituted by R⁸ and/or R⁹, in particular a 5-membered heteroaryl ring containing at least one heteroatom selected from oxygen, nitrogen and sulphur such as furyl, thienyl, isoxazolyl, imidazolyl or pyrazolyl optionally substituted by up to three R⁸ and/or R⁹ groups; and phenyl optionally substituted by R⁸ and/or R⁹.

Further representative examples of R⁵ include C₁₋₄alkyl, in particular methyl, phenyl optionally substituted by R⁸ and/or R⁹, and a 5-membered heteroaryl ring containing at least one heteroatom selected from oxygen, nitrogen and sulphur optionally substituted by R⁸ and/or R⁹, in particular thienyl optionally substituted by R⁸ and/or R⁹.

Representative examples of R⁶ include -(CH₂)₁heteroaryl optionally substituted by R¹¹ and/or R¹², in particular in a 5- or 6-membered heteroaryl containing at least one heteroatom selected from oxygen, nitrogen and sulfur, for example, pyridinyl optionally substituted by -(CH₂)₅NR¹⁶R¹⁷, furyl or thienyl.

Representative examples of R⁷ include C₃₋₇cycloalkyl, phenyl optionally substituted by R¹¹ and/or R¹², and heteroaryl optionally substituted by R¹¹ and/or R¹². In one embodiment, R⁷ is selected from C₃₋₆cycloalkyl such as cyclopropyl, cyclobutyl, cyclopentyl or cyclohexyl, in particular cyclopropyl; phenyl optionally substituted by C₁₋₄alkoxy, in particular methoxy, or -(CH₂)₅NR¹⁶R¹⁷; and heteroaryl optionally substituted by R¹¹ and/or R¹², in particular a 5- or 6-membered heteroaryl containing at least one heteroatom selected from oxygen, nitrogen and sulfur, for example, pyridinyl, thiazolyl or thiadiazolyl. In a further embodiment, R⁷ is C₃₋₆cycloalkyl, in particular cyclopropyl.

In one embodiment, R⁸ and R⁹ are each independently selected from halogen, cyano, trifluoromethyl, C₁₋₆alkyl, C₁₋₆alkoxy, -COR¹⁵, -CO₂R¹⁵, and heteroaryl.

Representative examples of R⁸ and R⁹ include halogen, in particular chlorine or fluorine; cyano; trifluoromethyl; nitro; C₁₋₄alkyl, in particular methyl, ethyl, n-propyl, isopropyl or n-butyl; C₁₋₄alkoxy, in particular methoxy; -CONR¹³R¹⁴; -COR¹⁵; -CO₂R¹⁵; and heteroaryl, in particular a 5-membered heteroaryl ring containing up to two heteroatoms independently selected from nitrogen and oxygen, for example isoxazolyl.

Further representative examples of R⁸ and R⁹ include halogen, in particular fluorine, cyano, trifluoromethyl, C₁₋₄alkoxy, in particular methoxy, COR¹⁵, CO₂R¹⁵, and heteroaryl, in particular a 5-membered heteroaryl ring containing up to two heteroatoms independently selected from nitrogen and oxygen, for example isoxazolyl.

In another embodiment, R⁸ and R⁹ are linked to form a fused 5-membered heterocyclyl ring containing one heteroatom selected from oxygen, sulphur and N-R¹⁰.

Representative examples include R⁸ and R⁹ being linked to form a fused 5-membered heterocyclyl ring containing oxygen or a fused 5-membered heteroaryl ring containing up to three heteroatoms selected from oxygen and nitrogen.

A further representative example is R⁸ and R⁹ being linked to form a fused 5-membered heterocyclyl ring containing oxygen.

Representative examples of R¹¹ and R¹² include C₁₋₄alkoxy, in particular methoxy, and -(CH₂)₅NR¹⁶R¹⁷.

Representative examples of R¹³ and R¹⁴ include hydrogen and C₁₋₄alkyl, in particular hydrogen and methyl.

Representative examples of R¹⁵ include C₁₋₄alkyl, in particular methyl.

In one embodiment R¹⁶ and R¹⁷, together with the nitrogen atom to which they are bound, form a 5- or 6-membered heterocyclic ring optionally further containing one additional oxygen atom, in particular pyrrolidinyl or morpholino.

In one embodiment, X and Y are each independently selected from hydrogen, chlorine and fluorine. Representative examples of X include hydrogen and fluorine. A representative example of Y is hydrogen. In a further embodiment, X and Y are each hydrogen.

In one embodiment, m is selected from 0, 1 and 2, in particular 0 and 1.

Representative examples of n include 0, 1 and 2. Further representative examples of n include 0 and 1.

Representative examples of q include 0 and 1, in particular 0.

A representative example of r is 0.

A representative example of s is 0.

It is to be understood that the present invention covers all combinations of particular and preferred groups described hereinabove. It is also to be understood that the present invention encompasses compounds of formula (I) in which a particular group or parameter, for example R¹⁰, R¹², R¹³, R¹⁴, R¹⁶, R¹⁷, m, q or s, may occur more than once. In such compounds it will be appreciated that each group or parameter is independently selected from the values listed.

Particular compounds according to the invention include those mentioned in the Examples. Specific examples which may be mentioned include:

N-[4-methyl-3-(3-piperidin-4-yl-1,2-benzisoxazol-6-yl)phenyl]-2-pyrrolidin-1-ylisonicotinamide;

N-cyclopropyl-4-methyl-3-[1-(phenylsulfonyl)-1H-indazol-5-yl]benzamide;

N-cyclopropyl-3-{1-[(3-fluorophenyl)sulfonyl]-1H-indazol-5-yl}-4-methylbenzamide;

N-cyclopropyl-4-methyl-3-[1-(methylsulfonyl)-1H-indazol-5-yl]benzamide;

N-cyclopropyl-3-{1-[(4-fluorophenyl)sulfonyl]-1H-indazol-5-yl}-4-methylbenzamide;

N-cyclopropyl-4-methyl-3-[3-(4-morpholinyl)-1,2-benzisoxazol-6-yl]benzamide;

- N-cyclopropyl-4-methyl-3-{3-[2-oxo-2-(1,3-thiazol-2-ylamino)ethyl]-1,2-benzisoxazol-6-yl}benzamide;
 N-cyclopropyl-4-methyl-3-[3-(4-morpholinylmethyl)-1,2-benzisoxazol-6-yl]benzamide;
 5 N-cyclopropyl-4-methyl-3-[3-(1-pyrrolidinylmethyl)-1,2-benzisoxazol-6-yl]benzamide;
 N-cyclopropyl-4-methyl-3-{1-[(1-methylethyl)sulfonyl]-1H-indazol-5-yl}benzamide);
 N-cyclopropyl-3-[1-(ethylsulfonyl)-1H-indazol-5-yl]-4-methylbenzamide;
 N-cyclopropyl-3-[1-(cyclopropylsulfonyl)-1H-indazol-5-yl]-4-methylbenzamide;
 10 N-cyclopropyl-4-methyl-3-[1-(2-thienylsulfonyl)-1H-indazol-5-yl]benzamide;
 3-{1-[(2-cyanophenyl)sulfonyl]-1H-indazol-5-yl}-N-cyclopropyl-4-methylbenzamide;
 N-cyclopropyl-3-fluoro-4-methyl-5-[1-(2-thienylsulfonyl)-1H-indazol-5-yl]benzamide;
 N-cyclopropyl-3-fluoro-4-methyl-5-[1-(3-thienylsulfonyl)-1H-indazol-5-yl]benzamide;
 15 N-cyclopropyl-3-fluoro-4-methyl-5-{1-[(1-methylethyl)sulfonyl]-1H-indazol-5-yl}benzamide;
 N-cyclopropyl-3-fluoro-4-methyl-5-[1-(propylsulfonyl)-1H-indazol-5-yl]benzamide;
 N-cyclopropyl-3-fluoro-4-methyl-5-[1-(methylsulfonyl)-1H-indazol-5-yl]benzamide;
 20 3-[1-(butylsulfonyl)-1H-indazol-5-yl]-N-cyclopropyl-5-fluoro-4-methylbenzamide;
 N-cyclopropyl-3-[1-(cyclopropylsulfonyl)-1H-indazol-5-yl]-5-fluoro-4-methylbenzamide;
 3-{1-[(5-chloro-2-thienyl)sulfonyl]-1H-indazol-5-yl}-N-cyclopropyl-5-fluoro-4-methylbenzamide;
 25 N-cyclopropyl-3-{1-[(3,5-dimethyl-4-isoxazolyl)sulfonyl]-1H-indazol-5-yl}-5-fluoro-4-methylbenzamide;
 N-cyclopropyl-3-[1-(ethylsulfonyl)-1H-indazol-5-yl]-5-fluoro-4-methylbenzamide;
 3-{1-[(2-chlorophenyl)sulfonyl]-1H-indazol-5-yl}-N-cyclopropyl-5-fluoro-4-methylbenzamide;
 30 N-cyclopropyl-3-fluoro-4-methyl-5-{1-[(2-methylphenyl)sulfonyl]-1H-indazol-5-yl}benzamide;
 N-cyclopropyl-3-fluoro-4-methyl-5-[1-(phenylsulfonyl)-1H-indazol-5-yl]benzamide;
 N-cyclopropyl-3-{1-[(2,5-difluorophenyl)sulfonyl]-1H-indazol-5-yl}-5-fluoro-4-methylbenzamide;
 35 N-cyclopropyl-3-fluoro-5-{1-[(2-fluorophenyl)sulfonyl]-1H-indazol-5-yl}-4-methylbenzamide;
 N-cyclopropyl-3-[1-(cyclopropylmethyl)-1H-indazol-5-yl]-4-methylbenzamide; and
 N-cyclopropyl-3-fluoro-4-methyl-5-[3-methyl-1-(2-thienylsulfonyl)-1H-indazol-5-yl]benzamide.

- 40 As used herein, the term "alkyl" refers to straight or branched hydrocarbon chains containing the specified number of carbon atoms. For example, C₁₋₆alkyl means a straight or branched alkyl containing at least 1, and at most 6, carbon atoms. Examples of "alkyl" as used herein include, but are not limited to, methyl, ethyl, n-

propyl, n-butyl, n-pentyl, isobutyl, isopropyl and t-butyl. A C₁₋₄alkyl group is preferred, for example methyl, ethyl, isopropyl or t-butyl. The said alkyl groups may be optionally substituted with one or more fluorine atoms for example, trifluoromethyl.

5 As used herein, the term "alkenyl" refers to straight or branched hydrocarbon chains containing the specified number of carbon atoms and containing at least one double bond. For example, C₂₋₆alkenyl means a straight or branched alkenyl containing at least 2, and at most 6, carbon atoms and containing at least one double bond. Examples of "alkenyl" as used herein include, but are not limited to ethenyl, 2-
10 propenyl, 3-butenyl, 2-butenyl, 2-pentenyl, 3-pentenyl, 3-methyl-2-butenyl, 3-methylbut-2-enyl, 3-hexenyl and 1,1-dimethylbut-2-enyl.

 As used herein, the term "alkoxy" refers to a straight or branched chain alkoxy group, for example, methoxy, ethoxy, propoxy, prop-2-oxy, butoxy, but-2-oxy, 2-methylprop-1-oxy, 2-methylprop-2-oxy, pentoxy, or hexyloxy. A C₁₋₄alkoxy group
15 is preferred, for example methoxy or ethoxy.

 As used herein, the term "cycloalkyl" refers to a non-aromatic hydrocarbon ring containing the specified number of carbon atoms which may optionally contain up to one double bond. For example, C₃₋₇cycloalkyl means a non-aromatic ring containing at least three, and at most seven, ring carbon atoms. Examples of
20 "cycloalkyl" as used herein include, but are not limited to, cyclopropyl, cyclobutyl, cyclopentyl, cyclohexyl and cycloheptyl. A C₃₋₆cycloalkyl group is preferred, for example, cyclopropyl, cyclopentyl or cyclohexyl.

 As used herein, the terms "heteroaryl ring" and "heteroaryl", unless otherwise defined, refer to a monocyclic 5- to 7-membered unsaturated hydrocarbon ring
25 containing at least one heteroatom independently selected from oxygen, nitrogen and sulfur. Preferably, the heteroaryl ring has five or six ring atoms. Examples of heteroaryl rings include, but are not limited to, furyl, thienyl, pyrrolyl, oxazolyl, thiazolyl, isoxazolyl, isothiazolyl, imidazolyl, pyrazolyl, oxadiazolyl, triazolyl, tetrazolyl, thiadiazolyl, pyridyl, pyridazinyl, pyrimidinyl, pyrazinyl and triazinyl. The
30 said ring may be optionally substituted by one or more substituents independently selected from C₁₋₆alkyl and oxy.

 As used herein, the terms "heterocyclic ring" or "heterocyclyl", unless otherwise defined refer to a monocyclic 3- to 7-membered saturated hydrocarbon ring containing at least one heteroatom independently selected from oxygen, nitrogen and
35 sulfur. Preferably, the heterocyclyl ring has five or six ring atoms. Examples of heterocyclyl groups include, but are not limited to, pyrrolidinyl, imidazolidinyl, pyrazolidinyl, piperidyl, piperazinyl, morpholino, tetrahydropyranyl, tetrahydrofuranyl, and thiomorpholino. The said ring may be optionally substituted by one or more substituents independently selected from C₁₋₆alkyl and oxy.

40 As used herein, the terms "halogen" or "halo" refer to the elements fluorine, chlorine, bromine and iodine. Preferred halogens are fluorine, chlorine and bromine. A particularly preferred halogen is fluorine or chlorine.

As used herein, the term "solvate" refers to a complex of variable stoichiometry formed by a solute (in this invention, a compound of formula (I) or a salt thereof) and a solvent. Such solvents for the purpose of the invention may not interfere with the biological activity of the solute. Examples of suitable solvents include water, methanol, ethanol and acetic acid. Preferably the solvent used is a pharmaceutically acceptable solvent. Examples of suitable pharmaceutically acceptable solvents include water, ethanol and acetic acid. All such solvates are included within the scope of the present invention.


Certain compounds of formula (I) may exist in stereoisomeric forms (e.g. they may contain one or more asymmetric carbon atoms or may exhibit cis-trans isomerism). The individual stereoisomers (enantiomers and diastereomers) and mixtures of these are included within the scope of the present invention. The present invention also covers the individual isomers of the compounds represented by formula (I) as mixtures with isomers thereof in which one or more chiral centres are inverted. Likewise, it is understood that compounds of formula (I) may exist in tautomeric forms other than that shown in the formula and these are also included within the scope of the present invention.

Salts of the compounds of the present invention are also encompassed within the scope of the invention and may, for example, comprise acid addition salts resulting from reaction of an acid with a basic nitrogen atom present in a compound of formula (I).

Salts encompassed within the term "pharmaceutically acceptable salts" refer to non-toxic salts of the compounds of this invention. Representative salts include the following salts: Acetate, Benzenesulfonate, Benzoate, Bicarbonate, Bisulphate, Bitartrate, Borate, Bromide, Calcium Edetate, Camsylate, Carbonate, Chloride, Clavulanate, Citrate, Dihydrochloride, Edetate, Edisylate, Estolate, Esylate, Fumarate, Gluceptate, Gluconate, Glutamate, Glycollylarsanilate, Hexylresorcinolate, Hydrabamine, Hydrobromide, Hydrochloride, Hydroxynaphthoate, Iodide, Isethionate, Lactate, Lactobionate, Laurate, Malate, Maleate, Mandelate, Mesylate, Methylbromide, Methylnitrate, Methylsulphate, Monopotassium Maleate, Mucate, Napsylate, Nitrate, N-methylglucamine, Oxalate, Pamoate (Embonate), Palmitate, Pantothenate, Phosphate/diphosphate, Polygalacturonate, Potassium, Salicylate, Sodium, Stearate, Subacetate, Succinate, Tannate, Tartrate, Teoclate, Tosylate, Triethiodide, Trimethylammonium and Valerate. Other salts which are not pharmaceutically acceptable may be useful in the preparation of compounds of this invention and these form a further aspect of the invention.


The compounds of this invention may be made by a variety of methods, including standard chemistry. Any previously defined variable will continue to have the previously defined meaning unless otherwise indicated. Illustrative general synthetic methods are set out below and then specific compounds of the invention are prepared in the working Examples.

A compound of formula (I) may be prepared by reacting a compound of formula (II)




(II)

in which A is as hereinbefore defined and Z¹ is halogen, in particular bromine,
 5 with a compound of formula (IIIA) or (IIIB)



(IIIA)




(IIIB)

10

in which R¹, R², X and Y are as hereinbefore defined,
 in the presence of a catalyst, for example tetrakis(triphenylphosphine)palladium.

A compound of formula (IIIA) may be prepared by, for example, reacting a
 15 compound of formula (IV)




(IV)

in which R¹, R², X and Y are as hereinbefore defined and Z² is halogen, in particular
 iodine,

with bis(pinnacolato)diboron, [1,1'-bis(diphenylphosphino)ferrocene] dichloropalladium (II) complex ($\text{PdCl}_2(\text{ppdf})$) and potassium acetate in a solvent such as DMF.

- 5 A compound of formula (IIIB) may be prepared by, for example, reacting a compound of formula (IV) as hereinbefore defined, with n-butyl lithium and triisopropyl borate in a solvent such as THF.

When R^2 is $-\text{NH}-\text{CO}-\text{R}^6$, a compound of formula (IV) may be prepared by reacting an amine of formula (V)



10

(V)

in which R^1 , X, Y and Z^2 are as hereinbefore defined, with an acid compound of formula (VI)

15




(VI)

in which R^6 is as hereinbefore defined, under amide forming conditions.

- 20 Suitable amide forming conditions are well known in the art and include adding a base such as DIPEA to a mixture of the amine of formula (V), the acid of formula (VI), and HATU in a solvent such as DMF.

Alternatively, when R^2 is $-\text{CO}-\text{NH}-(\text{CH}_2)_q-\text{R}^7$, a compound of formula (IV) may readily be prepared from a corresponding acid compound of formula (VII)




25

(VII)

in which R^1 , X, Y and Z^2 are as hereinbefore defined, by converting the acid to an activated form of the acid, for example the acid chloride, by treatment with, for example, thionyl chloride, and then reacting the activated acid

30 thus formed with an amine compound of formula (VIII)




(VIII)

in which R^7 is as hereinbefore defined,
under amide forming conditions.

Suitable amide forming conditions are well known in the art and include treating a solution of the acid of formula (VII), or the activated form thereof, in for
5 example DMF, with an amine of formula (VIII) in the presence of a base such as triethylamine.


For example, one general method for preparing the compounds of formula (I) comprises the reaction set out in Scheme 1 below.



Scheme 1


Alternatively, a further general method comprises final stage modification of one compound of formula (I) into another compound of formula (I). Suitable functional group transformations for converting one compound of formula (I) into another compound of formula (I) are well known in the art and are described in, for
15 instance, *Comprehensive Heterocyclic Chemistry II*, eds. A. R. Katritzky, C. W. Rees and E. F. V. Scriven (Pergamon Press, 1996), *Comprehensive Organic Functional Group Transformations*, eds. A.R. Katritzky, O. Meth-Cohn and C.W. Rees (Elsevier Science Ltd., Oxford, 1995), *Comprehensive Organic Chemistry*, eds. D. Barton and W.D. Ollis (Pergamon Press, Oxford, 1979), and *Comprehensive Organic*
20 *Transformations*, R.C. Larock (VCH Publishers Inc., New York, 1989).

For example, one method for preparing the compounds of formula (I) comprises the reactions set out in Scheme 2 below.




Scheme 2

For example, another method for preparing the compounds of formula (I) comprises the reactions set out in Scheme 3 below.



Scheme 3


For example, another method for preparing the compounds of formula (I) comprises the reactions set out in Scheme 4 below.



Scheme 4


For example, another method for preparing the compounds of formula (I)
5 comprises the reactions set out in Scheme 5 below.

16




Scheme 5

For example, another method for preparing the compounds of formula (I) comprises the reactions set out in Scheme 6 below.




Scheme 6

For example, another method for preparing the compounds of formula (I) comprises the reactions set out in Scheme 7 below.



Scheme 7

For example, another method for preparing the compounds of formula (I) comprises the reactions set out in Scheme 8 below.



Scheme 8

10

Those skilled in the art will appreciate that in the preparation of the compounds of the invention it may be necessary and/or desirable to protect one or

more sensitive groups in the molecule to prevent undesirable side reactions. Suitable protecting groups for use according to the present invention are well known to those skilled in the art and may be used in a conventional manner. See, for example, "Protective groups in organic synthesis" by T.W. Greene and P.G.M. Wuts (John Wiley & sons 1991) or "Protecting Groups" by P.J. Kocienski (Georg Thieme Verlag 1994). Examples of suitable amino protecting groups include acyl type protecting groups (e.g. formyl, trifluoroacetyl, acetyl), aromatic urethane type protecting groups (e.g. benzyloxycarbonyl (Cbz) and substituted Cbz), aliphatic urethane protecting groups (e.g. 9-fluorenylmethoxycarbonyl (Fmoc), t-butyloxycarbonyl (Boc), isopropylloxycarbonyl, cyclohexyloxycarbonyl) and alkyl type protecting groups (e.g. benzyl, trityl, chlorotriyl). Examples of suitable oxygen protecting groups may include for example alky silyl groups, such as trimethylsilyl or tert-butyldimethylsilyl; alkyl ethers such as tetrahydropyranyl or tert-butyl; or esters such as acetate.

Whilst it is possible for the compounds of the present invention to be administered as the new chemical, the compounds of formula (I) are conveniently administered in the form of pharmaceutical compositions. Thus, in another aspect of the invention, we provide a pharmaceutical composition comprising a compound of formula (I), in admixture with one or more pharmaceutically acceptable carriers, diluents or excipients.

The compounds of formula (I) may be formulated for administration in any suitable manner. They may, for example, be formulated for topical administration or administration by inhalation or, more preferably, for oral, transdermal or parenteral administration. The pharmaceutical composition may be in a form such that it can effect controlled release of the compounds of formula (I). A particularly preferred method of administration, and corresponding formulation, is oral administration.

For oral administration, the pharmaceutical composition may take the form of, and be administered as, for example, tablets (including sub-lingual tablets) and capsules (each including timed release and sustained release formulations), pills, powders, granules, elixirs, tinctures, emulsions, solutions, syrups or suspensions prepared by conventional means with acceptable excipients.

For instance, for oral administration in the form of a tablet or capsule, the active drug component can be combined with an oral, non-toxic pharmaceutically acceptable inert carrier such as ethanol, glycerol, water and the like. Powders are prepared by comminuting the compound to a suitable fine size and mixing with a similarly comminuted pharmaceutical carrier such as an edible carbohydrate, as, for example, starch or mannitol. Flavoring, preservative, dispersing and coloring agent can also be present.

Capsules can be made by preparing a powder mixture as described above, and filling formed gelatin sheaths. Glidants and lubricants such as colloidal silica, talc, magnesium stearate, calcium stearate or solid polyethylene glycol can be added to the powder mixture before the filling operation. A disintegrating or solubilizing agent such as agar-agar, calcium carbonate or sodium carbonate can also be added to improve the availability of the medicament when the capsule is ingested.

Moreover, when desired or necessary, suitable binders, lubricants, disintegrating agents and coloring agents can also be incorporated into the mixture. Suitable binders include starch, gelatin, natural sugars such as glucose or beta-lactose, corn sweeteners, natural and synthetic gums such as acacia, tragacanth or sodium alginate, carboxymethylcellulose, polyethylene glycol, waxes and the like. Lubricants used in these dosage forms include sodium oleate, sodium stearate, magnesium stearate, sodium benzoate, sodium acetate, sodium chloride and the like. Disintegrators include, without limitation, starch, methyl cellulose, agar, bentonite, xanthan gum and the like. Tablets are formulated, for example, by preparing a powder mixture, granulating or slugging, adding a lubricant and disintegrant and pressing into tablets. A powder mixture is prepared by mixing the compound, suitably comminuted, with a diluent or base as described above, and optionally, with a binder such as carboxymethylcellulose, an aliginat, gelatin, or polyvinyl pyrrolidone, a solution retardant such as paraffin, a resorption accelerator such as a quaternary salt and/or an absorption agent such as bentonite, kaolin or dicalcium phosphate. The powder mixture can be granulated by wetting with a binder such as syrup, starch paste, acadia mucilage or solutions of cellulosic or polymeric materials and forcing through a screen. As an alternative to granulating, the powder mixture can be run through the tablet machine and the result is imperfectly formed slugs broken into granules. The granules can be lubricated to prevent sticking to the tablet forming dies by means of the addition of stearic acid, a stearate salt, talc or mineral oil. The lubricated mixture is then compressed into tablets. The compounds of the present invention can also be combined with free flowing inert carrier and compressed into tablets directly without going through the granulating or slugging steps. A clear or opaque protective coating consisting of a sealing coat of shellac, a coating of sugar or polymeric material and a polish coating of wax can be provided. Dyestuffs can be added to these coatings to distinguish different unit dosages.

Oral fluids such as solution, syrups and elixirs can be prepared in dosage unit form so that a given quantity contains a predetermined amount of the compound. Syrups can be prepared by dissolving the compound in a suitably flavored aqueous solution, while elixirs are prepared through the use of a non-toxic alcoholic vehicle. Suspensions can be formulated by dispersing the compound in a non-toxic vehicle. Solubilizers and emulsifiers such as ethoxylated isostearyl alcohols and polyoxy ethylene sorbitol ethers, preservatives, flavor additives such as peppermint oil or saccharin, and the like can also be added.

Where appropriate, dosage unit formulations for oral administration can be microencapsulated. The formulation can also be prepared to prolong or sustain the release as for example by coating or embedding particulate material in polymers, wax or the like.

The compounds of the present invention can also be administered in the form of liposome delivery systems, such as small unilamellar vesicles, large unilamellar vesicles and multilamellar vesicles. Liposomes can be formed from a variety of phospholipids, such as cholesterol, stearylamine or phosphatidylcholines.

The compounds of the present invention can also be administered in the form of liposome emulsion delivery systems, such as small unilamellar vesicles, large unilamellar vesicles and multilamellar vesicles. Liposomes can be formed from a variety of phospholipids, such as cholesterol, stearylamine or phosphatidylcholines.

5 Compounds of the present invention may also be delivered by the use of monoclonal antibodies as individual carriers to which the compound molecules are coupled. The compounds of the present invention may also be coupled with soluble polymers as targetable drug carriers. Such polymers can include polyvinylpyrrolidone, pyran copolymer, polyhydroxypropylmethacrylamide-phenol, 10 polyhydroxyethylaspartamidephenol, or polyethyleneoxidepolylysine substituted with palmitoyl residues. Furthermore, the compounds of the present invention may be coupled to a class of biodegradable polymers useful in achieving controlled release of a drug, for example, polylactic acid, polepsilon caprolactone, polyhydroxy butyric acid, polyorthoesters, polyacetals, polydihydropyrans, polycyanoacrylates and cross-linked or amphipathic block copolymers of hydrogels. 15

The present invention includes pharmaceutical compositions containing 0.1 to 99.5%, more particularly, 0.5 to 90% of a compound of the formula (I) in combination with a pharmaceutically acceptable carrier.

20 Likewise, the composition may also be administered in nasal, ophthalmic, otic, rectal, topical, intravenous (both bolus and infusion), intraperitoneal, intraarticular, subcutaneous or intramuscular, inhalation or insufflation form, all using forms well known to those of ordinary skill in the pharmaceutical arts.

For transdermal administration, the pharmaceutical composition may be given in the form of a transdermal patch, such as a transdermal iontophoretic patch.

25 For parenteral administration, the pharmaceutical composition may be given as an injection or a continuous infusion (e.g. intravenously, intravascularly or subcutaneously). The compositions may take such forms as suspensions, solutions or emulsions in oily or aqueous vehicles and may contain formulatory agents such as suspending, stabilizing and/or dispersing agents. For administration by injection these 30 may take the form of a unit dose presentation or as a multidose presentation preferably with an added preservative. Alternatively for parenteral administration the active ingredient may be in powder form for reconstitution with a suitable vehicle.

35 The compounds of the invention may also be formulated as a depot preparation. Such long acting formulations may be administered by implantation (for example subcutaneously or intramuscularly) or by intramuscular injection. Thus, for example, the compounds of the invention may be formulated with suitable polymeric or hydrophobic materials (for example as an emulsion in an acceptable oil) or ion exchange resins, or as sparingly soluble derivatives, for example, as a sparingly soluble salt.

40 Alternatively the composition may be formulated for topical application, for example in the form of ointments, creams, lotions, eye ointments, eye drops, ear drops, mouthwash, impregnated dressings and sutures and aerosols, and may contain appropriate conventional additives, including, for example, preservatives, solvents to

assist drug penetration, and emollients in ointments and creams. Such topical formulations may also contain compatible conventional carriers, for example cream or ointment bases, and ethanol or oleyl alcohol for lotions. Such carriers may constitute from about 1% to about 98% by weight of the formulation; more usually they will
5 constitute up to about 80% by weight of the formulation.

For administration by inhalation the compounds according to the invention are conveniently delivered in the form of an aerosol spray presentation from pressurized packs or a nebulizer, with the use of a suitable propellant, e.g. dichlorodifluoromethane, trichlorofluoromethane, dichlorotetrafluoroethane,
10 tetrafluoroethane, heptafluoropropane, carbon dioxide or other suitable gas. In the case of a pressurized aerosol the dosage unit may be determined by providing a valve to deliver a metered amount. Capsules and cartridges of e.g. gelatin for use in an inhaler or insufflator may be formulated containing a powder mix of a compound of the invention and a suitable powder base such as lactose or starch.

15 The pharmaceutical compositions generally are administered in an amount effective for treatment or prophylaxis of a specific condition or conditions. Initial dosing in human is accompanied by clinical monitoring of symptoms, such symptoms for the selected condition. In general, the compositions are administered in an amount of active agent of at least about 100 $\mu\text{g/kg}$ body weight. In most cases they will be administered in
20 one or more doses in an amount not in excess of about 20 mg/kg body weight per day. Preferably, in most cases, dose is from about 100 $\mu\text{g/kg}$ to about 5 mg/kg body weight, daily. For administration particularly to mammals, and particularly humans, it is expected that the daily dosage level of the active agent will be from 0.1 mg/kg to 10 mg/kg and typically around 1 mg/kg. It will be appreciated that optimum dosage will be
25 determined by standard methods for each treatment modality and indication, taking into account the indication, its severity, route of administration, complicating conditions and the like. The physician in any event will determine the actual dosage which will be most suitable for an individual and will vary with the age, weight and response of the particular individual. The effectiveness of a selected actual dose can readily be
30 determined, for example, by measuring clinical symptoms or standard anti-inflammatory indicia after administration of the selected dose. The above dosages are exemplary of the average case. There can, of course, be individual instances where higher or lower dosage ranges are merited, and such are within the scope of this invention. For conditions or disease states as are treated by the present invention, maintaining
35 consistent daily levels in a subject over an extended period of time, e.g., in a maintenance regime, can be particularly beneficial.

In another aspect, the present invention provides a compound of formula (I) for use in therapy.

40 The compounds of the present invention are generally inhibitors of the serine/threonine kinase p38 and are therefore also inhibitors of cytokine production which is mediated by p38 kinase. Within the meaning of the term "inhibitors of the serine/threonine kinase p38" are included those compounds that interfere with the

ability of p38 to transfer a phosphate group from ATP to a protein substrate according to the assay described below.

It will be appreciated that the compounds of the invention may be selective for one or more of the isoforms of p38, for example p38 α , p38 β , p38 γ and/or p38 δ . In one embodiment, the compounds of the invention selectively inhibit the p38 α isoform. In another embodiment, the compounds of the invention selectively inhibit the p38 β isoform. In a further embodiment, the compounds of the invention selectively inhibit the p38 α and p38 β isoforms. Assays for determining the selectivity of compounds for the p38 isoforms are described in, for example, WO 99/61426, WO 00/71535 and WO 02/46158.

It is known that p38 kinase activity can be elevated (locally or throughout the body), p38 kinase can be incorrectly temporally active or expressed, p38 kinase can be expressed or active in an inappropriate location, p38 kinase can be constitutively expressed, or p38 kinase expression can be erratic; similarly, cytokine production mediated by p38 kinase activity can be occurring at inappropriate times, inappropriate locations, or it can occur at detrimentally high levels.

Accordingly, the present invention provides a method for the treatment of a condition or disease state mediated by p38 kinase activity, or mediated by cytokines produced by the activity of p38 kinase, in a subject which comprises administering to said subject a therapeutically effective amount of a compound of formula (I). The compound may be administered as a single or polymorphic crystalline form or forms, an amorphous form, a single enantiomer, a racemic mixture, a single stereoisomer, a mixture of stereoisomers, a single diastereoisomer or a mixture of diastereoisomers.

The present invention also provides a method of inhibiting cytokine production which is mediated by p38 kinase activity in a subject, e.g. a human, which comprises administering to said subject in need of cytokine production inhibition a therapeutic, or cytokine-inhibiting, amount of a compound of the present invention. The compound may be administered as a single or polymorphic crystalline form or forms, an amorphous form, a single enantiomer, a racemic mixture, a single stereoisomer, a mixture of stereoisomers, a single diastereoisomer or a mixture of diastereoisomers.

The present invention treats these conditions by providing a therapeutically effective amount of a compound of this invention. By "therapeutically effective amount" is meant a symptom-alleviating or symptom-reducing amount, a cytokine-reducing amount, a cytokine-inhibiting amount, a kinase-regulating amount and/or a kinase-inhibiting amount of a compound. Such amounts can be readily determined by standard methods, such as by measuring cytokine levels or observing alleviation of clinical symptoms. For example, the clinician can monitor accepted measurement scores for anti-inflammatory treatments.

The compounds of the present invention can be administered to any subject in need of inhibition or regulation of p38 kinase or in need of inhibition or regulation of p38 mediated cytokine production. In particular, the compounds may be administered to mammals. Such mammals can include, for example, horses, cows, sheep, pigs,

mice, dogs, cats, primates such as chimpanzees, gorillas, rhesus monkeys, and, most preferably, humans.

Thus, the present invention provides methods of treating or reducing symptoms in a human or animal subject suffering from, for example, rheumatoid arthritis, osteoarthritis, asthma, psoriasis, eczema, allergic rhinitis, allergic conjunctivitis, adult respiratory distress syndrome, chronic pulmonary inflammation, chronic obstructive pulmonary disease, chronic heart failure, silicosis, endotoxemia, toxic shock syndrome, inflammatory bowel disease, tuberculosis, atherosclerosis, neurodegenerative disease, Alzheimer's disease, Parkinson's disease, Huntington's disease, amyotrophic lateral sclerosis, epilepsy, multiple sclerosis, aneurism, stroke, irritable bowel syndrome, muscle degeneration, bone resorption diseases, osteoporosis, diabetes, reperfusion injury, graft vs. host reaction, allograft rejections, sepsis, systemic cachexia, cachexia secondary to infection or malignancy, cachexia secondary to acquired immune deficiency syndrome (AIDS), malaria, leprosy, infectious arthritis, leishmaniasis, Lyme disease, glomerulonephritis, gout, psoriatic arthritis, Reiter's syndrome, traumatic arthritis, rubella arthritis, Crohn's disease, ulcerative colitis, acute synovitis, gouty arthritis, spondylitis, and non articular inflammatory conditions, for example, herniated/ruptured/prolapsed intervertebral disk syndrome, bursitis, tendonitis, tenosynovitis, fibromyalgic syndrome and other inflammatory conditions associated with ligamentous sprain and regional musculoskeletal strain, pain, for example that associated with inflammation and/or trauma, osteopetrosis, restenosis, thrombosis, angiogenesis, cancer including breast cancer, colon cancer, lung cancer or prostatic cancer, which comprises administering to said subject a therapeutically effective amount of a compound of formula (I).

A further aspect of the invention provides a method of treatment of a human or animal subject suffering from rheumatoid arthritis, asthma, psoriasis, chronic pulmonary inflammation, chronic obstructive pulmonary disease, chronic heart failure, systemic cachexia, glomerulonephritis, Crohn's disease, neurodegenerative disease, Alzheimer's disease, Parkinson's disease, epilepsy and cancer including breast cancer, colon cancer, lung cancer and prostatic cancer, which comprises administering to said subject a therapeutically effective amount of a compound of formula (I).

A further aspect of the invention provides a method of treatment of a human or animal subject suffering from rheumatoid arthritis, asthma, psoriasis, chronic pulmonary inflammation, chronic obstructive pulmonary disease, chronic heart failure, systemic cachexia, glomerulonephritis, Crohn's disease and cancer including breast cancer, colon cancer, lung cancer and prostatic cancer, which comprises administering to said subject a therapeutically effective amount of a compound of formula (I).

A further aspect of the invention provides a method of treatment of a human or animal subject suffering from rheumatoid arthritis, asthma, chronic pulmonary inflammation, chronic obstructive pulmonary disease, neurodegenerative disease,

Alzheimer's disease, Parkinson's disease and epilepsy which comprises administering to said subject a therapeutically effective amount of a compound of formula (I).

5 A further aspect of the invention provides a method of treatment of a human or animal subject suffering from any type of pain including chronic pain, rapid onset of analgesis, neuromuscular pain, headache, cancer pain, acute and chronic inflammatory pain associated with osteoarthritis and rheumatoid arthritis, post operative
10 inflammatory pain, neuropathic pain, diabetic neuropathy, trigeminal neuralgia, post-hepatic neuralgia, inflammatory neuropathies and migraine pain which comprises administering to said subject a therapeutically effective amount of a compound of formula (I) or a pharmaceutically acceptable salt or solvate thereof.

The compounds of formula (I) may be employed alone or in combination with other therapeutic agents for the treatment of the above-mentioned conditions. In particular, in rheumatoid arthritis therapy, combination with other chemotherapeutic or antibody agents is envisaged. Combination therapies according to the present
15 invention thus comprise the administration of at least one compound of formula (I) and at least one other pharmaceutically active agent. The compound(s) of formula (I) and the other pharmaceutically active agent(s) may be administered together or separately and, when administered separately, this may occur separately or sequentially in any order. The amounts of the compound(s) of formula (I) and the
20 other pharmaceutically active agent(s) and the relative timings of administration will be selected in order to achieve the desired combined therapeutic effect. Examples of other pharmaceutically active agents which may be employed in combination with compounds of formula (I) for rheumatoid arthritis therapy include:
25 immunosuppressants such as amtolmetin guacil, mizoribine and rimexolone; anti-TNF α agents such as etanercept, infliximab, diacerein; tyrosine kinase inhibitors such as leflunomide; kallikrein antagonists such as subreum; interleukin 11 agonists such as oprelvekin; interferon beta 1 agonists; hyaluronic acid agonists such as NRD-101 (Aventis); interleukin 1 receptor antagonists such as anakinra; CD8 antagonists such as amiprilose hydrochloride; beta amyloid precursor protein antagonists such as
30 reumacon; matrix metalloprotease inhibitors such as cipemastat and other disease modifying anti-rheumatic drugs (DMARDs) such as methotrexate, sulphasalazine, cyclosporin A, hydroxychloroquine, auranofin, aurothioglucose, gold sodium thiomalate and penicillamine.

Examples

The following Examples are illustrative embodiments of the invention, not limiting the scope of the invention in any way. Reagents are commercially available or are prepared according to procedures in the literature.

LCMS was conducted on a column (3.3cm x 4.6mm ID, 3µm ABZ+PLUS), at a Flow Rate of 3ml/min, Injection Volume of 5µl, at room temperature and UV Detection Range at 215 to 330nm.

2-Chloroisonicotinic acid, 3-cyanobenzenesulfonyl chloride, 3,4-difluorobenzenesulfonyl chloride, 2,3-dihydro-1-benzofuran-5-sulfonyl chloride and 5-isoxazol-3-ylthiophene-2-sulfonyl chloride may be purchased from Maybridge Chemicals.

3-Iodo-4-methylaniline, 3-fluorobenzenesulfonyl chloride and 4-(trifluoromethyl)benzenesulfonyl chloride may be purchased from Avocado.

3-Methoxyphenylsulfonyl chloride may be purchased from Lancaster.

4-Acetylphenylsulfonyl chloride may be purchased from Acros.


5-Bromoindazole may be prepared by the procedure described in Chem Ber, 1922, 55, 1141.

6-Bromo-3-piperidin-4-yl-1,2-benzisoxazole may be prepared by the procedure described in WO97/49698.

6-Bromo-3-methyl-1,2-benzisoxazole and 5-bromo-3-methyl-1,2-benzisoxazole may be prepared according to the procedures described in Indian J Chem Sect B, 1977, 15B, 1058-1063.

Methyl 4-[(chlorosulfonyl)methyl]benzoate may be prepared by the procedure described in Rec Trav Chim Pays-Bas, 1957, 76, 129.


Intermediate 1: 6-Bromo-3-bromomethyl-1,2-benzisoxazole



A mixture of 6-bromo-3-methyl-1,2-benzisoxazole (675mg), N-bromosuccinimide (642mg) and 1,1'-azobis(cyclohexanecarbonitrile) (90mg) in carbon tetrachloride (7ml) was irradiated using a 300W lamp at reflux under nitrogen for 40h. On cooling the precipitate was removed by filtration, the solvent was evaporated and the residue was purified by flash chromatography on a silica column (4cm diam) eluting with a cyclohexane-ethyl acetate gradient (49:1 to 24:1) to give the title compound as a pale yellow solid (350mg).

LC-MS: Rt 3.67min.

Intermediate 2: 2-[(6-Bromo-1,2-benzisoxazol-3-yl)methyl]amino-1,3-propanediol



5


A solution of 6-bromo-3-bromomethyl-1,2-benzisoxazole (Intermediate 1, 130mg), serinol (50mg) and DIPEA (0.05ml) in DMF (0.2ml) was stirred at 60° under nitrogen for 3h. The mixture was partitioned between water and ethyl acetate and the organic layer was washed with water and brine, dried through a hydrophobic filter and concentrated. The residue was purified on a Varian Bond Elut SPE cartridge (silica, 2g) eluting with dichloromethane followed by dichloromethane-methanol (9:1) to give the title compound as a yellow gum (33mg).

10

LC-MS: Rt 1.75min, MH⁺ 301,303.

15

Intermediate 3: tert-Butyl 5-bromo-1H-indazole-1-carboxylate



20

A stirred ice-cold suspension of 5-bromoindazole (2g, 10.2mmol), 4-(dimethylamino)pyridine (250mg, 2.0mmol) and triethylamine (1.55ml, 11.2mmol) in acetonitrile (50ml) was treated with a solution of di-tert-butyl dicarbonate (2.8ml, 12.2mmol) in acetonitrile (20ml) over 15 min such that the temperature remained under 5°. The reaction mixture was warmed to room temp then stirred for 18h. The solvent was evaporated and the residue was purified by column chromatography on silica (100g) eluting with cyclohexane:ethyl acetate (15:1) to give the title compound (2.27g, 7.7mmol).


25

NMR: δ H[CDCl₃] 8.10 (1H, s), 8.07 (1H, d), 7.86 (1H, d), 7.60 (1H, dd), 1.71 (9H, s).

30

LCMS: Rt 3.55min.

Intermediate 4: 4-Methyl-3-(4,4,5,5-tetramethyl-[1,3,2]dioxaborolan-2-yl)benzoic acid



5


3-Iodo-4-methylbenzoic acid (10g, 38.16mmol), bis(pinnacolato)diboron (14.5g, 57.24mmol), potassium acetate (18.73g, 190.8mmol) and PdCl₂dppf (3.12g, 3.8mmol) in DMF (200ml) were heated at 80°C for 21hrs. The solvent was evaporated from the cooled reaction under vacuum, the residue dissolved in ethyl acetate (300ml) and hydrochloric acid (2N, 300ml) and filtered through celite. The organic phase was separated and the aqueous extracted with ethyl acetate (2 x 300ml). The combined organic extracts were washed with brine (500ml) and dried (magnesium sulphate). The solvent was evaporated under vacuum and the residue was absorbed onto silica and applied to a silica column. This was eluted with cyclohexane / ethyl acetate (5:1). to give the title compound.

NMR: δ H [d6-DMSO] 12.83,(1H, b), 8.23,(1H, d), 7.89,(1H, dd), 7.29,(1H, d), 2.51,(3H, s), 1.30,(12H, s).

LCMS: Rt 3.65min.

20

Intermediate 5: N-Cyclopropyl-4-methyl-3-(4,4,5,5-tetramethyl-1,3,2-dioxaborolan-2-yl)benzamide




Intermediate 4 (10g) was dissolved in DMF (100ml). To this was added cyclopropylamine (2.2g) DIPEA (15ml) and HATU (14g). The mixture was stirred for 3 hours at room temp. The solvent was removed under vacuum and the residue was partitioned between ethyl acetate (400ml) and saturated sodium bicarbonate solution (400ml). The organic layer was dried over magnesium sulphate, filtered and concentrated under vacuum to give a pink solid. The product was purified using

25

Biotage chromatography on silica eluting with 50:50 ethylacetate/cyclohexane to give the title compound as a white solid.


- 5 NMR: δ H [d6-DMSO] 8.42 (1H, d), 8.06 (1H, d), 7.77 (1H, dd), 7.23 (1H, d), 2.85 (1H, m), 2.47 (3H, s), 1.32 (12H, s), 0.67 (2H, m), 0.56 (2H, m).
LCMS: Rt 3.29min, MH^+ 302.

Intermediate 6: 3-Iodo-4-methylbenzoyl chloride




- 10 Thionyl chloride (8.2ml) was added to a mixture of 3-iodo-4-methylbenzoic acid (18.5g) in chloroform (100ml) and heated at 61°C for 16 hours. The solvent was removed under vacuum and excess thionyl chloride removed by azeotrope with toluene (3x30ml). The title compound was formed as a beige solid (19.5g) and used in subsequent reactions without further purification.
- 15 NMR: δ H [d6-DMSO] 8.31 (1H, d), 7.87 (1H, dd), 7.46 (1H, d), 2.43 (3H, s).

Intermediate 7: 4-(3-Nitrophenyl)morpholine



- 20 3-Fluoronitrobenzene (10g) was added to a solution of morpholine (34ml) in DMSO (120ml) and heated at 110°C for 60h. The reaction mixture was cooled and poured onto water (800ml). The precipitate was collected by filtration and the orange solid was dried under vacuum and used in subsequent reactions without further purification (13.7g).
- 25 NMR: δ H [d6-DMSO] 7.68 (1H, dd), 7.62 (1H, dd), 7.49 (1H, t), 7.42 (1H, dd), 3.76 (4H, dd), 3.24 (4H, dd).


Intermediate 8: 3-(4-Morpholinyl)benzenamine



A flask containing 5% palladium on carbon (1.95g) was evacuated and refilled with hydrogen. 4-(3-Nitrophenyl)morpholine (Intermediate 7) (19.5g) was introduced into the flask as a solution in ethanol and DMF (1000ml, 4:1 v/v). The reaction was stirred at room temp until further uptake of hydrogen ceased (after approximately 7L). The reaction was then filtered through celite and solvent was removed under vacuum to give the title compound (12.6g) as a beige solid.

NMR: δ H [d6-DMSO] 6.85 (1H, t), 6.12 (2H, m), 6.06 (1H, dd), 4.88 (2H, brs), 3.70 (4H, apparent t), 2.98 (4H, apparent t).
LCMS: Rt 1.08 min MH^+ 179.


Intermediate 9: 3-Iodo-4-methyl-N-[3-(4-morpholinyl)phenyl]benzamide



3-Iodo-4-methylbenzoyl chloride (Intermediate 6) (19.5g) was added portion-wise to a mixture of triethylamine (48ml) and 3-(4-morpholinyl)benzenamine (Intermediate 8) (12.6g) in DMF (150ml) and the mixture was heated at 80°C for 16h. The solvent was removed under vacuum and the residue was dissolved in chloroform (200ml). The organic layer was washed with water (2x100ml), 2M sodium hydroxide solution (100ml) and brine (100ml), dried over magnesium sulphate, filtered and concentrated under vacuum. The resulting yellow solid was triturated with diethyl ether and collected by filtration to yield the title compound as an off-white solid (20.0g). The product was used in subsequent reactions without further purification.

NMR: δ H [d6-DMSO] 10.10 (1H, s), 8.39 (1H, d), 7.90 (1H, dd), 7.49 (1H, d), 7.38 (1H, t), 7.28 (1H, brd), 7.19 (1H, t), 6.71 (1H, dd), 3.75 (4H, apparent t), 3.10 (4H, apparent t), 2.44 (3H, s).
LCMS: Rt 3.52 min MH^+ 423.


Intermediate 10: 4-Methyl-N-[3-(4-morpholinyl)phenyl]-3-(4,4,5,5-tetramethyl-[1,3,2]-dioxaborolan-2-yl)benzamide



3-Iodo-4-methyl-N-[3-(4-morpholinyl)phenyl]benzamide (Intermediate 9) (8.00g)
5 triethylamine (7.9ml) and bis(pinacolato)diboron (4.13ml) were added to a solution of PdCl₂dppf (770mg) in dioxane (100ml) and the mixture was heated at 80° under nitrogen for 3h. The reaction was cooled, the solvent was removed under vacuum and the residue was dissolved in dichloromethane (150ml). The solution was washed with water (100mlx3) and brine (100ml) dried (magnesium sulphate) and the solvent was
10 removed under vacuum. The residue was purified by column chromatography on silica eluting with 30% ethyl acetate/cyclohexane to 50% ethyl acetate/cyclohexane to give the title compound as a white solid (4.05g).

NMR: δ H [d₆-DMSO] 10.11 (1H, s), 8.19 (1H, d), 7.93 (1H, dd), 7.40 (1H, brs), 7.33
15 (1H, d), 7.28 (1H, brd), 7.19 (1H, t), 6.70 (1H, dd), 3.75 (4H, apparent t), 3.09 (4H, apparent t), 2.54 (3H, s), 1.33 (12H, s).
LCMS: Rt 3.65 min MH⁺423.

Intermediate 11: 2-Chloro-N-(3-iodo-4-methylphenyl)-isonicotinamide




20 2-Chloroisonicotinic acid (3.3g) HATU (8.75g) DIPEA (10.9ml) and 3-iodo-4-methylaniline (5.00g) in DMF (50ml) were heated under nitrogen for 16h. The reaction was cooled, the solvent was removed under vacuum and the residue was dissolved in dichloromethane (150ml). The solution was washed with water
25 (3x100ml) and brine (100ml) dried (magnesium sulphate) and the solvent was removed under vacuum. The residue was purified by column chromatography on silica eluting with ethyl acetate/cyclohexane (40:60) to give the title compound as a white solid (7.00g).

NMR: δ H [d6-DMSO] 10.52 (1H, s), 8.62 (1H, d), 8.29 (1H, d), 7.99 (1H, b), 7.87 (1H, dd), 7.70 (1H, dd), 7.34 (1H, d), 2.36 (3H, s).

LCMS: Rt 3.59 min MH^+ 373.

5 **Intermediate 12: N-(3-Iodo-4-methylphenyl)-2-pyrrolidin-1-yl-isonicotinamide**




10 A solution of N-(3-iodo-4-methylphenyl)-2-chloro-isonicotinamide (Intermediate 11) (7.00g) in pyrrolidine (20ml) was heated at 80°C under nitrogen for 16h. Excess pyrrolidine was removed under vacuum and the residue was triturated with diethyl ether (20ml). The resulting solid was collected by filtration and dried under vacuum to give the title compound as a pale yellow solid (7.73g).

15 NMR: δ H [d6-DMSO] 10.29 (1H, s), 8.29 (1H, d), 8.20 (1H, d), 7.71 (1H, dd), 7.72 (1H, dd), 6.97 (1H, brd), 6.88 (1H, b), 3.45 (2H, apparent t), 3.09 (2H, m), 2.35 (3H, s), 1.98 (2H, m), 1.82 (2H, m).

LCMS: Rt 2.77 min MH^+ 408.

Intermediate 13: N-[4-Methyl-3-(4,4,5,5-tetramethyl-[1,3,2]-dioxaborolan-2-yl)phenyl]-2-(1-pyrrolidinyl)-4-pyridinecarboxamide




20 Bis(pinacolato)diborane (7.24g) was added to a mixture of N-(3-iodo-4-methylphenyl)-2-pyrrolidin-1-yl-isonicotinamide (Intermediate 12) (7.73g) potassium acetate (9.32g, 95mmol) and $PdCl_2dppf$ (770mg) in DMF (100ml) and the reaction was heated at 80° under nitrogen for 16h. The reaction was cooled and the solvent was removed under vacuum. The residue was dissolved in chloroform (150ml), washed
25 with water (3x100ml) and brine (100ml), dried (magnesium sulphate) and the solvent was removed under vacuum. The residue was purified by column chromatography on

silica eluting with ethyl acetate:cyclohexane (1:4) to (1:1) to give the title compound as a white solid (1.5g).

5 NMR: δ H (CDCl₃) 8.27 (1H, d), 7.99 (1H, dd), 7.76 (1H, b), 7.65 (1H, d), 6.20 (1H, d), 6.82 (1H, b), 6.77 (1H, b), 3.52 (4H, apparent t), 2.52 (3H, s), 2.25 (4H, m), 1.35 (12H, s).

LCMS: Rt 2.90 min MH⁺408.

Intermediate 14: N-(3-Iodo-4-methylphenyl)thiophene-3-amide



10

Thiophene-3-carboxylic acid (2.75g) and HATU (8.15g) in DMF (25ml) were stirred at room temp for 15min. HOBT (2.9g), 3-iodo-4-methylaniline (5.0g) and DIPEA (11.2ml, 64.35mmol) were added and the reaction was stirred at room temp for 16h. The solvent was evaporated under vacuum and the residue was partitioned between ethyl acetate (100ml) and aqueous sodium carbonate (10%, 100ml). The aqueous layer was extracted with ethyl acetate (50ml) and the combined organic phases washed with hydrochloric acid (2N, 75ml), water (75ml) and brine (75ml). The organic phase was dried (magnesium sulphate) and absorbed onto silica. The silica was applied to a silica column and eluted with cyclohexane/ethyl acetate (4:1). The solvent was evaporated from the product fractions under vacuum to the title compound.


15

20

NMR: δ H [d6-DMSO] 10.06,(1H, b), 8.34,(1H, m), 8.29,(1H, d), 7.70,(1H, dd), 7.66,(1H, dd), 7.62,(1H, dd), 7.30,(1H, d), 2.34,(3H, s).

25 LCMS: Rt 3.69min, MH⁺ 344.

Intermediate 15: N-[4-Methyl-3-(4,4,5,5-tetramethyl-[1,3,2]dioxaborolan-2-yl)-phenyl]thiophene-3-amide



30


N-(3-Iodo-4-methylphenyl)thiophene-3-amide (Intermediate 14) (2.64g) bis(pinacolato)diboron (2.13g), potassium acetate (825mg) and PdCl₂dppf (312mg)

in DMF (20ml) were heated at 80°C for 20h. The cooled reaction was absorbed onto silica and applied to a Varian Bond-Elut SPE cartridge (silica, 10g) and eluted with a cyclohexane/ethyl acetate gradient. The product fractions were concentrated under vacuum to give the title compound.

5

NMR: δ H [d6-DMSO] 9.99,(1H, b), 8.35,(1H, s), 7.90,(1H, d), 7.85,(1H, dd), 7.63,(2H, m), 7.14,(1H, d), 2.42,(3H, s), 1.30,(12H, s).
LCMS: Rt 3.65min, MH^+ 344.


10 **Intermediate 16: N-(3-Iodo-4-methylphenyl)-3-furamide**



3-Furoic acid (2.4g) and HATU (8.15g) in DMF (25ml) were stirred at room temperature for 15min. HOBT (2.9g), 3-iodo-4-methylaniline (5.0g) and DIPEA (11.2ml) were added and the mixture was stirred at room temp for 16h. The solvent was evaporated under vacuum and the residue was partitioned between ethyl acetate (100ml) and aqueous sodium carbonate (10%, 100ml). The aqueous layer was extracted with ethyl acetate (50ml) and the combined organic phases were washed with hydrochloric acid (2N, 75ml), water (75ml) and brine (75ml). The organic phase was dried (magnesium sulphate) and absorbed onto silica. The silica was applied to a silica column and eluted with cyclohexane/ethyl acetate (3:1) to give the title compound.

NMR: δ H [d6-DMSO] 9.92,(1H, b), 8.36,(1H, d), 8.23,(1H, d), 7.80,(1H, t), 7.66,(1H, dd), 7.29,(1H, d), 6.98,(1H, d), 2.33,(3H, s).
LCMS: Rt 3.52min, MH^+ 328.

Intermediate 17: N-[4-Methyl-3-(4,4,5,5-tetramethyl-[1,3,2]dioxaborolan-2-yl)-phenyl]-3-furamide



30 N-(3-Iodo-4-methylphenyl)-3-furamide (Intermediate 16) (2.5g) bis(pinacolato)diboron (2.13g) potassium acetate (825mg) and $PdCl_2dppf$ (312mg) in


DMF (20ml) were heated at 80° for 20h. The cooled reaction was absorbed onto silica and applied to a Varian Bond-Elut SPE cartridge (silica, 10g) and eluted with a cyclohexane/ethyl acetate gradient. The product fractions were concentrated under vacuum to give the title compound.

5

NMR: δ H [d6-DMSO] 9.86,(1H, b), 8.36,(1H, m), 7.86-7.82,(2H, m), 7.77,(1H, t), 7.14,(1H, d), 6.99,(1H, m), 2.41,(3H, s), 1.30,(12H, s).

LCMS: Rt 3.55min, MH^+ 328.

10 **Intermediate 18: 4-Methyl-3-(4,4,5,5-tetramethyl-[1,3,2]dioxaborolan-2-yl)-N-(thiazol-2-yl)-benzamide**




15 4-Methyl-3-(4,4,5,5-tetramethyl-[1,3,2]dioxaborolan-2-yl)-benzoic acid (Intermediate 4) (2.0g) DIPEA (4ml) and HATU (3.05g) were dissolved in DMF (20ml) and stirred at room temp for 15min. 2-Aminothiazole (801mg) was added and the mixture was stirred at room temp for 18h. The solvent was evaporated under vacuum and the residue was partitioned between ethyl acetate (250ml) and water (50ml). The organic phase was washed with hydrochloric acid (2N, 50ml) and aqueous sodium bicarbonate (1M, 50ml) dried (magnesium sulphate) concentrated under vacuum. The residue was
20 absorbed onto silica and purified by flash column chromatography eluting with cyclohexane/ethyl acetate (4:1) to give the title compound (1.72g).

NMR: δ H [d6-DMSO] 12.65,(1H, b), 8.32,(1H, d), 8.08,(1H, dd), 7.56,(1H, d), 7.35,(1H, d), 7.28,(1H, d), 2.54,(3H, s), 1.34,(12H, s).

25 LCMS: Rt 3.66min, MH^+ 345.

Intermediate 19: N-(3-Methoxy-phenyl)-4-methyl-3-(4,4,5,5-tetramethyl-[1,3,2]dioxaborolan-2-yl)-benzamide




- 4-Methyl-3-(4,4,5,5-tetramethyl-[1,3,2]dioxaborolan-2-yl)-benzoic acid (Intermediate 4) (2g) was dissolved in DMF (20ml). To this was added 3-methoxyaniline (0.985g), DIPEA (4ml) and HATU (3.05g). The mixture was stirred for 18 h at room temp. The solvent was removed under vacuum and the residue was partitioned between ethyl acetate (250ml) and water (50ml). The organic layer was dried (magnesium sulphate) concentrated under vacuum and purified using a silica Biotage cartridge (90g) eluting with 1:4 ethyl acetate/cyclohexane to give the title compound as a white solid (2.06g).

NMR: δ H [d6-DMSO] 10.20, (1H, s), 8.17, (1H, s), 7.94-7.91, (1H, dd), 7.45, (1H, s), 7.36-7.32, (2H, t), 7.25-7.21, (1H, t), 6.68-6.65, (1H, dd), 3.74, (3H, s), 2.53, (3H, s), 1.32, (12H, s).

- LCMS: Rt 3.80min, MH^+ 368.

Intermediate 20: 4-Methyl-3-(4,4,5,5-tetramethyl-[1,3,2]dioxaborolan-2-yl)-N-([1,3,4]thiadiazol-2-yl)-benzamide




- 4-Methyl-3-(4,4,5,5-tetramethyl-[1,3,2]dioxaborolan-2-yl)-benzoic acid (Intermediate 4) (2.0g), DIPEA (4ml) and HATU (3.05g) were dissolved in DMF (20ml) and stirred at room temp for 15min. 2-Aminothiadiazole (810mg) was added and the mixture was stirred at room temp for 18h. The solvent was evaporated under vacuum and the residue was partitioned between ethyl acetate (250ml) and hydrochloric acid (2N, 150ml). The aqueous phase was extracted with ethyl acetate (2 x 250ml). The combined organic extracts were dried (magnesium sulphate) and the solvent was removed under vacuum. The residue was absorbed onto silica and purified by flash column chromatography eluting with cyclohexane/ethyl acetate (4:1 then 1:1) to give the title compound (0.95g).

NMR: δ H [d6-DMSO] 13.08,(1H, b), 9.22,(1H, s), 8.35,(1H, d), 8.11,(1H, dd), 7.38,(1H, d), 2.55,(3H, s), 1.34,(12H, s).

LCMS: Rt 3.34min, MH^+ 346.

5

Intermediate 21: N-Cyclopropylmethyl-4-methyl-3-(4,4,5,5-tetramethyl-[1,3,2]dioxaborolan-2-yl)-benzamide




10 4-Methyl-3-(4,4,5,5-tetramethyl-[1,3,2]dioxaborolan-2-yl)-benzoic acid (Intermediate 4) (2.0g) DIPEA (4ml) and HATU (3.05g) were dissolved in DMF (20ml) and stirred at room temp for 15min. Cyclopropylmethylamine (568mg) was added and the mixture was stirred at room temp for 18h. The solvent was removed under vacuum and the residue was partitioned between ethyl acetate (250ml) and water (50ml). The organic phase was washed with hydrochloric acid (2N, 50ml) and aqueous sodium bicarbonate (1M, 50ml) dried (magnesium sulphate) and concentrated under vacuum.

15 The residue was absorbed onto silica and purified by flash column chromatography eluting with cyclohexane / ethyl acetate (4:1) to give the title compound (1.73g).

20 NMR: δ H [d6-DMSO] 8.54,(1H, t), 8.11,(1H, d), 7.82,(1H, dd), 7.26,(1H, d), 3.12,(2H, t), 1.32,(12H, s), 1.03,(1H, m), 0.42,(2H, m), 0.22,(2H, m).

LCMS: Rt 3.47min, MH^+ 316.

Intermediate 22: Methyl 4-bromo-2-hydroxybenzoate



25


A solution of 4-bromo-2-hydroxybenzoic acid (1.0g) in methanol (10ml) was treated cautiously with conc. sulphuric acid (0.5ml) then stirred at 75° under nitrogen for 7h. The mixture was concentrated under vacuum and the residue was partitioned between ethyl acetate (30ml) and water (30ml). The aqueous layer was re-extracted with ethyl acetate (30ml) and the combined organic extracts were washed with saturated aqueous sodium bicarbonate (2x50ml) and water (50ml). The dried ($MgSO_4$) extracts were concentrated under vacuum to give the title compound as a beige solid (0.7g).

30

LCMS: Rt 3.42min.

Intermediate 23: 4-Bromo-N,2-dihydroxybenzamide


5



10 A solution of methyl 4-bromo-2-hydroxybenzoate (Intermediate 22) (0.7g) in dioxan (5ml) was added dropwise to a solution of hydroxylamine hydrochloride (0.32g) and sodium hydroxide (0.42g) in water (10ml) then stirred at room temp. for 18h. The dioxan was removed under vacuum and the residue was stirred with 2N hydrochloric acid to give a precipitate which was collected by filtration, washed with water and dried to give the title compound as an off-white solid (0.63g).

15 LCMS: Rt 2.64min.

Intermediate 24: 6-Bromo-1,2-benzisoxazol-3(2H)-one




20

25 A solution of carbonyldiimidazole (0.75g) in dry tetrahydrofuran (15ml) was added to a refluxing solution of 4-bromo-N,2-dihydroxybenzamide (Intermediate 23) (0.56g) in dry tetrahydrofuran (10ml) then stirred for 2h. The solvent was removed under vacuum, the residue was suspended in water and 2N hydrochloric acid (10ml) was added. The resulting precipitate was collected by filtration, washed with water then cyclohexane and dried to give the title compound as a cream solid (0.37g).

LCMS: Rt 3.30min.

30 **Intermediate 25: 1,1-Dimethylethyl 4-(6-bromo-1,2-benzisoxazol-3-yl)-1-piperazinecarboxylate**




5 An ice-bath cooled solution of 6-bromo-1,2-benzisoxazol-3(2H)-one (Intermediate 24) (0.05g) and dry pyridine (0.05ml) in dry dichloromethane (4ml) was stirred under nitrogen and treated with triflic anhydride (60 μ L). The solution was stirred at room temp. for 4h, diluted with cyclohexane then applied to a Varian Bond-Elut SPE
10 cartridge (silica, 1g) and eluted with dichloromethane to give a yellow oil. The oil was dissolved in acetonitrile (2ml) treated with 1,1-dimethylethyl 1-piperazinecarboxylate (0.04g) and diisopropylethylamine (0.05ml) then stirred at 70° under nitrogen for 20h. The cooled reaction mixture was purified on a Varian Bond-Elut SPE cartridge (silica, 5g) using dichloromethane:methanol to give the title compound as a white solid (0.016g).

LCMS: Rt 3.62min.

15

Intermediate 26: 1,1-Dimethylethyl 4-(6-{5-[(cyclopropylamino)carbonyl]-2-methylphenyl}-1,2-benzisoxazol-3-yl)-1-piperazinecarboxylate




20

A mixture of 1,1-dimethylethyl 4-(6-bromo-1,2-benzisoxazol-3-yl)-1-piperazinecarboxylate (Intermediate 25) (0.04g), *N*-Cyclopropyl-4-methyl-3-(4,4,5,5-tetramethyl-1,3,2-dioxaborolan-2-yl)benzamide (Intermediate 5) (0.035g), 2N aqueous sodium carbonate (2ml) and tetrakis(triphenylphosphine)palladium(0) (1mg) in isopropanol (6ml) was stirred at reflux under nitrogen for 16h. The residue was
25 absorbed onto silica (Merck 7734) and applied to a Varian Bond-Elut SPE cartridge (silica, 5g). Elution with dichloromethane:methanol (98:2) gave the title compound as a cream solid (0.03g).

LCMS: Rt 3.50min.

Intermediate 27: 6-Bromo-3-(4-morpholinyl)-1,2-benzisoxazole

5




10 An ice-bath cooled solution of 6-bromo-1,2-benzisoxazol-3(2H)-one (Intermediate 24) (0.15g) and dry pyridine (0.15ml) in dry dichloromethane (12ml) was stirred under nitrogen and treated with triflic anhydride (180μL). The solution was stirred at room temp. for 90min, diluted with cyclohexane then applied to a Varian Bond-Elut SPE cartridge (silica, 5g) and eluted with cyclohexane and dichloromethane to give a colourless oil. The oil was dissolved in acetonitrile (4ml) treated with morpholine (61 μL) and diisopropylethylamine (0.15ml) then stirred at 70° under nitrogen for 16h. The cooled reaction mixture was purified on a Varian Bond-Elut SPE cartridge (silica, 5g) using dichloromethane:ethanol:0.88ammonia to give the title compound as a white solid (0.03g).

LCMS: Rt 3.01min.

20

Intermediate 28: 3-Bromophenyl acetate




25 Acetic anhydride was added dropwise to an ice-cold solution of 3-bromophenol (25.5g) in pyridine (25ml) and the solution was then stirred at room temp. for 4h. 2N Hydrochloric acid (160ml) was added and the mixture was extracted with ether (x2). The combined organic extracts were washed with water and brine, dried (MgSO₄) and concentrated under vacuum to give the title compound as a brown liquid (30.8g).

30

LCMS: Rt 3.0min.

Intermediate 29: 1-(4-Bromo-2-hydroxyphenyl)ethanone




3-Bromophenyl acetate (Intermediate 28) (30.6g) was added portionwise to powdered aluminium chloride 15.5g) with stirring. The resulting liquid was stirred at 170° under nitrogen for 1h then at 70° for 2h. 2N Hydrochloric acid (150ml) was added and the mixture was allowed to cool to room temp. The mixture was extracted with ethyl acetate (x2) and the solvent was evaporated to leave a dark oil which was purified on silica (500g) eluting with cyclohexane:ethyl acetate (4:1) to give the title compound as a slightly coloured a semi-solid (26.8g).

10

LCMS: Rt 3.2min.

Intermediate 30: Ethyl 3-(4-bromo-2-hydroxyphenyl)-3-oxopropanoate



15


A solution of ethyl 3-(4-bromo-2-hydroxyphenyl)-3-oxopropanoate (Intermediate 29) (3.0g) in dry toluene (10ml) was added dropwise to a stirred suspension of sodium hydride (60% oil dispersion, 1.54g) in dry toluene (15ml) at reflux under nitrogen. After the addition stirring was continued for a further 10min then diethylcarbonate (3.39ml) was added over 30 min. The mixture was then stirred at reflux for 18h. On cooling, ice-cold 2N hydrochloric (150ml) added and the mixture was extracted with ethyl acetate (2x100ml). The combined extracts were washed with water and brine, dried (MgSO₄) and concentrated under vacuum. The residue was purified by column chromatography on silica (100g) eluting with cyclohexane:ethylacetate (10:1) to give the title compound as a cream solid (2.87g).

20

25

LCMS: Rt 3.25min.

30 **Intermediate 31: 7-Bromo-4-hydroxy-2H-chromen-2-one**




A solution of ethyl 3-(4-bromo-2-hydroxyphenyl)-3-oxopropanoate (Intermediate 30) (2.0g) in dry toluene (20ml) was stirred at reflux under nitrogen for 24h. On cooling the resulting precipitate was collected by filtration, washed with ether and dried to give the title compound as a white solid (1.32g).

5

LCMS: Rt 3.47min.

Intermediate 32: (6-Bromo-1,2-benzisoxazol-3-yl)acetic acid

10



15


Sodium metal (0.36g) was dissolved in dry ethanol (30ml) then treated with hydroxylamine hydrochloride (1.12g). A solution of 7-bromo-4-hydroxy-2H-chromen-2-one (Intermediate 31) (1.3g) in dry ethanol (30ml) was added and the mixture was stirred at reflux under nitrogen for 3.5h. On cooling the mixture was added to 2M aqueous sodium bicarbonate (100ml). Water (100ml) and dichloromethane (100ml) were added, the phases were separated and the aqueous layer was washed with more dichloromethane (75ml). The aqueous layer was acidified by the cautious addition of 2N hydrochloric acid to yield a precipitate which was collected by filtration and dried to give the title compound as a white solid (1.06g).

20

LCMS: Rt 2.99min.

Intermediate 33: 1,1-Dimethylethyl 4-[(6-bromo-1,2-benzisoxazol-3-yl)acetyl]-1-piperazinecarboxylate

25



30

A suspension of (6-bromo-1,2-benzisoxazol-3-yl)acetic acid (Intermediate 32) (0.06g) and HATU (0.09g) in dry tetrahydrofuran (2ml) was stirred at room temp. for 5min. 1-Hydroxybenzotriazole (0.03g) and 1,1-dimethylethyl 1-piperazinecarboxylate (0.04g) were added and the reaction mixture was stirred for a further 18h. The solvent was removed under vacuum and the residue was partitioned between dichloromethane and 2N aqueous sodium carbonate. The layers were separated and the organic layer was applied to a Varian Bond-Elut SPE cartridge (silica, 5g) eluting sequentially with


35

dichloromethane, chloroform, ether and ethyl acetate to give the title compound as a white solid (0.09g).

LCMS: Rt 3.21min.

5

Intermediate 34: 1,1-Dimethylethyl 4-[(6-{5-[(cyclopropylamino)carbonyl]-2-methylphenyl}-1,2-benzisoxazol-3-yl)acetyl]-1-piperazinecarboxylate



10

A mixture of 1,1-dimethylethyl 4-[(6-bromo-1,2-benzisoxazol-3-yl)acetyl]-1-piperazinecarboxylate (Intermediate 33) (0.045g) *N*-cyclopropyl-4-methyl-3-(4,4,5,5-tetramethyl-1,3,2-dioxaborolan-2-yl)benzamide (Intermediate 5) (0.03g) and 2M aqueous sodium carbonate (0.65ml) in isopropanol (2.5ml) under nitrogen was treated with tetrakis(triphenylphosphine)palladium(0) (1mg) then stirred at reflux for 18h. On cooling the mixture was absorbed onto silica (Merck 7734) and applied to a Varian Bond-Elut SPE cartridge (silica, 5g). Sequential elution with dichloromethane, chloroform, ether, ethyl acetate, acetonitrile and acetone yielded the title compound as a white solid (0.03g).


15

20

LCMS: Rt 3.16min.

Intermediate 35: 2-(6-Bromo-1,2-benzisoxazol-3-yl)-N-(2-hydroxyethyl)acetamide

25




30

A suspension of (6-bromo-1,2-benzisoxazol-3-yl)acetic acid (Intermediate 32) (0.10g) and HATU (0.18g) in tetrahydrofuran (5ml) was stirred at room temp. for 10 min then treated with 1-hydroxybenzotriazole (0.05g) 2-aminoethanol (0.03ml) and diisopropylethylamine (0.2ml). The mixture was stirred for a further 18h then concentrated under vacuum. The residue was partitioned between dichloromethane (5ml) and 2M aqueous sodium carbonate (5ml). The organic layer was dried using a

hydrophobic filter tube then applied to a Varian Bond-Elut SPE cartridge. Sequential elution with dichloromethane, ether, ethyl acetate, acetonitrile and acetone yielded the title compound as a white solid (0.08g).

5 LCMS: Rt 2.42min.

Intermediate 36: 6-Bromo-3-[2-oxo-2-(1-piperidinyl)ethyl]-1,2-benzisoxazole




10

The procedure for Intermediate 35 was followed using piperidine (0.05ml) in place of 2-aminoethanol to give the title compound as a white solid (0.10g).

LCMS: Rt 3.09min.

15

Intermediate 37: 2-(6-Bromo-1,2-benzisoxazol-3-yl)-N-methylacetamide




20 The procedure for Intermediate 35 was followed using a solution of methylamine in tetrahydrofuran (2M, 0.235ml) in place of 2-aminoethanol to give the title compound as a white solid (0.10g).

LCMS: Rt 2.59min.

25

Intermediate 38: 2-(6-Bromo-1,2-benzisoxazol-3-yl)-N-(3-hydroxypropyl)acetamide




30

The procedure for Intermediate 35 was followed using 3-aminopropanol (0.04ml) in place of 2-aminoethanol to give the title compound as a white solid (0.09g).

LCMS: Rt 2.48min.

Intermediate 39: 2-(6-Bromo-1,2-benzisoxazol-3-yl)-N-(cyclopropylmethyl)acetamide

5




The procedure for Intermediate 35 was followed using cyclopropylmethylamine (0.04ml) in place of 2-aminoethanol to give the title compound as a white solid (0.11g).

10

LCMS: Rt 2.99min.

Intermediate 40: 6-Bromo-3-[2-oxo-2-(1-pyrrolidinyl)ethyl]-1,2-benzisoxazole

15




The procedure for Intermediate 35 was followed using pyrrolidine (0.04ml) in place of 2-aminoethanol to give the title compound as a white solid (0.10g).

20

LCMS: Rt 2.87min.

Intermediate 41: 2-(6-Bromo-1,2-benzisoxazol-3-yl)-N-ethylacetamide

25




The procedure for Intermediate 35 was followed using a solution of ethylamine in tetrahydrofuran (0.235ml) in place of 2-aminoethanol to give the title compound as a white solid (0.09g).

30

LCMS: Rt 2.74min.

Intermediate 42: 2-(6-Bromo-1,2-benzisoxazol-3-yl)-N-cyclopropylacetamide




The procedure for Intermediate 35 was followed cyclopropylamine (0.03ml) in place of 2-aminoethanol to give the title compound as a white solid (0.08g).

5

LCMS: Rt 2.78min.

Intermediate 43: 6-Bromo-3-[2-(4-morpholinyl)-2-oxoethyl]-1,2-benzisoxazole




10

The procedure for Intermediate 35 was followed morpholine (0.04ml) in place of 2-aminoethanol to give the title compound as a white solid (0.10g).

15 LCMS: Rt 2.69min.

Intermediate 44: 2-(6-Bromo-1,2-benzisoxazol-3-yl)-N-([3-(methoxy)phenyl]methyl)acetamide




20

The procedure for Intermediate 35 was followed 3-methoxybenzylamine (0.06ml) in place of 2-aminoethanol to give the title compound as a white solid (0.10g).

25 LCMS: Rt 3.20min.


Intermediate 45: 2-(6-Bromo-1,2-benzisoxazol-3-yl)-N-1,3-thiazol-2-ylacetamide



The procedure for Intermediate 35 was followed 2-amino-1,3-thiazole (0.05ml) in place of 2-aminoethanol to give the title compound as a white solid (0.06g).

5 LCMS: Rt 3.11min.

Intermediate 46: 6-Bromo-3-[(4-methyl-1-piperazinyl)methyl]-1,2-benzisoxazole



10


A solution of Intermediate 1 (0.06g), 1-methylpiperazine (0.05ml) and diisopropylethylamine (0.1ml) in DMF (0.5ml) was stirred at 65° for 4h. The reaction mixture was partitioned between ethyl acetate and saturated aqueous sodium hydrogen carbonate, the phases were separated and the aqueous layer was re-extracted with ethyl acetate. The organic extracts were washed with water (x2) and brine, dried using a hydrophobic filter tube and concentrated under a stream of nitrogen. The residue was purified on a Varian Bond-Elut SPE cartridge (silica, 2g) eluting with dichloromethane:methanol:ethylamine (100:0:1 to 95:5:1) to give the title compound as a pale yellow solid (0.04g).

20

LCMS: Rt 2.07min.

Intermediate 47: 6-Bromo-3-(1-piperidinylmethyl)-1,2-benzisoxazole

25




The procedure for Intermediate 46 was followed using piperidine (0.05ml) in place of 1-methylpiperazine. Elution of the SPE cartridge with cyclohexane:ethyl acetate:triethylamine (100:0:1 to 80:20:1) gave the title compound as a pale yellow solid (0.03g).

30

LCMS: Rt 2.00min.

Intermediate 48: 6-Bromo-3-(4-morpholinylmethyl)-1,2-benzisoxazole


35



The procedure for Intermediate 46 was followed using morpholine (0.05ml) in place of 1-methylpiperazine. Elution of the SPE cartridge with cyclohexane:ethyl acetate:triethylamine (100:0:1 to 66:33:1) gave the title compound as a pale yellow solid (0.04g).

LCMS: Rt 2.46min.


Intermediate 49: 6-Bromo-3-(1-pyrrolidin-2-ylmethyl)-1,2-benzisoxazole



The procedure for Intermediate 46 was followed using pyrrolidine (0.05ml) in place of 1-methylpiperazine. Elution of the SPE cartridge with cyclohexane:ethyl acetate:triethylamine (100:0:1 to 80:20:1) gave the title compound as a pale yellow solid (0.03g).


LCMS: Rt 1.88min.

Intermediate 50: 4-Bromo-2-fluorobenzaldehyde oxime




A solution of hydroxylamine in water (50%, 5ml) was added to a solution of 4-bromo-2-fluorobenzaldehyde (1.0g) in ethanol (5ml) and the mixture was stirred at room temp. for 66h. The ethanol was removed under vacuum and the aqueous residue as extracted with ethyl acetate (x2). The organic extracts were washed with water and brine, dried (MgSO₄) and concentrated under vacuum. The residue was purified on a Varian Bond-Elut SPE cartridge (silica, 20g) eluting with cyclohexane:ethyl acetate (100:0 to 50:50) to give the title compound as a white solid (0.94g).

LCMS: Rt 3.15min.

Intermediate 51: 4-Bromo-2-fluoro-N-hydroxybenzenecarboximidoyl chloride

5 N-Chlorosuccinimide (0.075g) was added to a solution of 4-bromo-2-fluorobenzaldehyde oxime (Intermediate 50) (0.44g) in DMF (1ml) then heated for several minutes at 55°. The mixture was allowed to cool to <50° then more N-chlorosuccinimide (0.22g) was added. After a few minutes the mixture was allowed to
10 cool to room temp. then partitioned between water and ether. The layers were separated, the aqueous layer was re-extracted with ether (x2) and the combined organic extracts were washed with water and brine, dried (MgSO₄) and concentrated under vacuum to give a white solid (0.46g).

15 LCMS: Rt 3.11.


Intermediate 52: 4-Bromo-N-cyclopropyl-2-fluoro-N'-hydroxybenzenecarboximidamide

20

Cyclopropylamine (1.6ml) was added dropwise to an ice-cold solution of 4-bromo-2-fluoro-N-hydroxybenzenecarboximidoyl chloride (Intermediate 51) (0.59g) in dry ether (25ml) under nitrogen. After the addition the mixture was allowed to warm to
25 room temp. then stirred for 18h. The resulting precipitate was collected by filtration then purified by column chromatography on silica eluting with cyclohexane:ethyl acetate (3:2) to give the title compound as a white solid (0.435g).

LCMS: Rt 2.69min.


30

Intermediate 53: 6-Bromo-N-cyclopropyl-1,2-benzisoxazol-3-amine

1,8-Diazabicyclo[5.4.0]undec-7-ene (0.12ml) was added to 4-bromo-N-cyclopropyl-2-fluoro-N'-hydroxybenzenecarboximidamide (Intermediate 52) (0.2g) in tetrahydrofuran (2ml) then heated at 150° in a microwave for 70min. The mixture was applied to a Varian Bond-Elut SPE cartridge (silica, 20g) then eluted with cyclohexane:ethyl acetate (100:0 to 80:20) to give the title compound as a white solid (0.13g).

LCMS: Rt 3.16min.


Intermediate 54: Methyl (4-bromo-2-nitrophenyl)acetate



Conc. hydrochloric acid (1ml) was added to a solution of (4-bromo-2-nitrophenyl)acetic acid (7.07g) and the solution was stirred under reflux for 5h. The solvent was evaporated to leave the title compound as a crystalline brown solid (7.39g).

LCMS: Rt 3.13min.


Intermediate 55: Ethyl 6-bromo-1,2-benzisoxazole-3-carboxylate



Sodium metal (0.09g) was dissolved in absolute ethanol (2ml) under nitrogen. A solution of methyl (4-bromo-2-nitrophenyl)acetate (Intermediate 54) (1.0g) in ethanol (8ml) was treated with isoamyl nitrite (0.6ml) followed by the solution of sodium ethoxide, yielding a black mixture. The mixture was stirred at 60° for 90min then left at room temp. for 18h. 2N Hydrochloric acid (20ml) was added and the mixture was extracted with ethyl acetate (2x20ml). The combined organic extracts were washed with water (x2) and brine, dried using a hydrophobic filter tube and concentrated under vacuum. The residue was purified on a Varian Bond-Elut SPE cartridge eluting with cyclohexane:ethyl acetate (100:0 to 97:3) to give the title compound as a cream coloured solid (577mg).


LCMS: Rt 3.40min.

5 **Intermediate 56: 6-Bromo-N-(cyclopropylmethyl)-1,2-benzisoxazole-3-carboxamide**



10 A solution of ethyl 6-bromo-1,2-benzisoxazole-3-carboxylate (Intermediate 55) (0.05g) and cyclopropylamine (0.03ml) in methanol (1ml) was stirred at reflux for 6h. The solvent was evaporated and the residue was triturated with ether to give the title compound as a white solid (0.04g).


15 **Intermediate 57: 6-Bromo-N-propyl-1,2-benzisoxazole-3-carboxamide**



20 The procedure for Intermediate 56 was followed using propylamine (0.025ml) in place of cyclopropylmethylamine to give the title compound as a white solid (0.04g).

LCMS: Rt 3.30min.

Intermediate 58: 6-Bromo-N-methyl-1,2-benzisoxazole-3-carboxamide




25

The procedure for Intermediate 56 was followed using a solution of methylamine in methanol (2M, 0.45ml) in place of cyclopropylmethylamine to give the title compound as a white solid (0.03g).

5 LCMS: Rt 2.92min.

Intermediate 59: 6-Bromo-N,N-dimethyl-1,2-benzisoxazole-3-carboxamide




10

The procedure for Intermediate 56 was followed using a solution of dimethylamine in methanol (2M, 0.13ml) in place of cyclopropylmethylamine to give the title compound as a white solid (0.02g).

15 LCMS: Rt 2.88min.

Intermediate 60: 6-Bromo-N-cyclopropyl-1,2-benzisoxazole-3-carboxamide




20

The procedure for Intermediate 56 was followed using cyclopropylamine (0.025ml) in place of cyclopropylmethylamine to give the title compound as a white solid (0.05g).

LCMS: Rt 3.30min.

25


Intermediate 61: 3-Bromo-N-cyclopropyl-5-fluoro-4-methylbenzamide



3-Fluoro-4-methylbenzoic acid (0.46g) was added to a stirred mixture of bromine (2.31ml) and iron powder (0.25g) under nitrogen. The reaction was stirred at 20°C for 4h and then left to stand for 16h. Sodium thiosulphate solution (200ml) was added and the product was extracted into ethyl acetate (3 x 150ml). Ethyl acetate extracts were combined and concentrated under vacuum. The crude product was dissolved in DMF (7ml) and cyclopropylamine (0.28ml) 1-hydroxybenzotriazole (0.405g) 1-(3-dimethylaminopropyl)-3-ethylcarbodiimide hydrochloride (0.575g) and diisopropylethylamine (0.52ml) were added. The mixture was stirred for 5h at 20°C. The solvent was removed under vacuum and the residue was partitioned between ethyl acetate and water. The combined organic extracts were washed with aqueous sodium hydrogen carbonate and hydrochloric acid (0.5M) dried (MgSO₄) and concentrated under vacuum. The residue was purified by column chromatography on silica eluting with cyclohexane:ethyl acetate (6:1) to give the title compound (359mg).

LCMS: MH⁺ 272.


Intermediate 62: N-Cyclopropyl-3-fluoro-4-methyl-5-(4,4,5,5-tetramethyl-1,3,2-dioxaborolan-2-yl)benzamide



3-Bromo-N-cyclopropyl-5-fluoro-4-methylbenzamide (Intermediate 61) (900mg), bispinnacolatodiboron (4.5g), potassium acetate (2.1g) and PdCl₂dppf (75mg) were mixed in DMF (40ml) and heated at 100°C for 18h. The cooled reaction was absorbed onto silica and applied to a Varian Bond Elut SPE cartridge (silica, 2 x 10g) and eluted with an cyclohexane:ethyl acetate (100:0 to 94:6). The solvent was under vacuum and the residue was recrystallised from cyclohexane to give the title compound (260mg).

LCMS: Rt 3.39mins.


Intermediate 63: N-Cyclopropyl-3-fluoro-5-iodo-4-methylbenzamide



5 N-Iodosuccinimide (22.5g) was added in portions to a solution of 3-fluoro-4-methylbenzoic acid (15.4g) in trifluoromethanesulphonic acid (100ml) at 0°C over 3h and the mixture was allowed to warm to room temp overnight. The reaction mixture was poured into ice/water (400ml) and the precipitate was collected by filtration and washed with water. The solid was dissolved in ethyl acetate, washed with aqueous sodium thiosulphate (x2) and brine, dried (MgSO₄) and the solvent was removed under vacuum. The residue was treated with thionyl chloride (30ml) and heated at 100°C for 2.5h. Excess thionyl chloride was removed under vacuum and the residue was dissolved in dichloromethane (100ml). Sodium carbonate (25g) and cyclopropylamine (13ml) were added to the solution and the mixture was stirred at room temp. for 72h. The mixture was filtered and the residue was washed with dichloromethane and ethyl acetate. The combined filtrate and washings were concentrated under vacuum. The residue was purified by column chromatography on silica eluting with cyclohexane:ethyl acetate (78:22 to 72:28) to give the title compound.

LCMS: Rt 3.16min.

Intermediate 64: {5-[(Cyclopropylamino)carbonyl]-3-fluoro-2-methylphenyl}boronic acid




25 N-Cyclopropyl-5-fluoro-3-iodo-4-methylbenzamide (Intermediate 63) (5g) in tetrahydrofuran (75ml) was cooled to 0°C then treated with sodium hydride (60% oil dispersion, 1.23g) portionwise over 10minutes. Once effervescence had ceased the reaction was cooled to -75°C and n-butyl lithium (1.6M in hexanes, 20ml) was added over 25min maintaining a temperature of <-70°C. Triisopropyl borate (8ml) was added over 10min and the mixture was stirred at -70°C for 4h. Water (20ml) was added and the mixture was allowed to warm to 5°C. The solvent was removed under vacuum and the residue was partitioned between saturated aqueous ammonium chloride and ethyl acetate. The organic phase was washed with saturated aqueous ammonium chloride and brine, dried (Na₂SO₄) and concentrated under vacuum. The residue was dissolved in dichloromethane/ethyl acetate and purified by column

chromatography on silica eluting with dichloromethane:ethyl acetate (95:5 to 0:100) then methanol to give the title compound.

LCMS: Rt 2.19min.

5


Intermediate 65: 1-Acetyl-6-bromo-1H-indazole



- 10 A solution of 4-bromo-2-methylaniline (10.0g) in chloroform (100ml) was treated with acetic anhydride (11.5g) and potassium acetate (5.8g) then stirred at 55° for 5h to give a thick suspension. t-Butyl nitrite (19.2ml) and 18-crown-6 (28.4g) were added to give a solution which was stirred at 55° for a further 21h. The solvent was evaporated and the oily residue which was eluted through a pad of silica (150g) using
- 15 cyclohexane:ethyl acetate (4:1). The solvent was evaporated and the residue was purified by column chromatography on silica eluting with cyclohexane:ethyl acetate (20:1 to 8:1) to give the title compound (5.08g).

LCMS: Rt 3.20min.


20 **Intermediate 66: (4-Bromo-2-nitrophenyl)methyl 4-fluorophenyl sulfide**



- 25 To a solution of 4-bromo-1-(bromomethyl)-2-nitrobenzene (0.60g) in tetrahydrofuran (10ml) was added 4-fluorothiophenol (1.2ml) and diisopropylethylamine (0.45ml). After stirring for 2h the mixture was concentrated under vacuum and purified by column chromatography on silica, eluting with petroleum ether (40-60°):dichloromethane (9:1 to 8:1) to give the title compound (0.52g).

30 LCMS: Rt 3.73 min.


Intermediate 67: (4-Bromo-2-nitrophenyl)methyl 4-fluorophenyl sulfone



A suspension of oxone (3.55g) in water (8ml) was added to a solution of (4-bromo-2-nitrophenyl)methyl 4-fluorophenyl sulfide (Intermediate 66) (0.52g) in acetonitrile (10ml). The mixture was stirred for 5h filtered and the residue was washed with ethyl acetate. The biphasic filtrate was separated, the aqueous layer was re-extracted with ethyl acetate (2 x 50ml) and the combined organic extracts were washed with aqueous sodium metabisulphite (5%, 100ml) and brine. The dried (MgSO₄) extracts were concentrated under vacuum to give the title compound (0.56g).

LCMS: Rt 3.33min.


Intermediate 68: (5-Bromo-2-[(4-fluorophenyl)sulfonyl]methyl)phenylamine



Saturated ammonium chloride (2.2ml) and iron powder (0.25g) was added to a suspension of (5-bromo-2-[(4-fluorophenyl)sulfonyl]methyl)phenylamine (Intermediate 67) (0.56g) in toluene (10ml) then stirred for 18h at reflux. The mixture was cooled and evaporated to give a white solid which was partitioned between ethyl acetate (3 x 20ml) and saturated aqueous sodium bicarbonate (15ml). The dried (MgSO₄) organic extracts were concentrated under vacuum to give the title compound (0.42g).

LCMS: Rt 3.16 min.

Intermediate 69: 6-Bromo-3-[(4-fluorophenyl)sulfonyl]-1H-indazole




To a suspension of (5-bromo-2-((4-fluorophenyl)sulfonyl)methylphenyl)amine (Intermediate 68) (0.42g) in chloroform (10ml) was added potassium acetate (0.135g) and acetic anhydride (0.24ml). The mixture was heated under nitrogen at reflux for 18h. 18-Crown-6 (0.64g) and tert-butylnitrite (0.435ml) were added and the mixture was stirred at reflux for a further 42h. The mixture was concentrated under vacuum and the residue was purified by column chromatography on silica eluting with cyclohexane:ethylacetate (4:1) to give the title compound (0.16g).

10

LCMS: Rt 3.43 min.

Intermediate 70: 6-Bromo-3-[(4-fluorophenyl)sulfonyl]-1-([2-(trimethylsilyl)ethyl]oxy)methyl]-1H-indazole



15


Sodium tert-butoxide (0.035g) was added to an ice-cold solution of 6-bromo-3-[(4-fluorophenyl)sulfonyl]-1H-indazole (Intermediate 69) (0.11g) in tetrahydrofuran (1.5ml). 2-(Trimethylsilyl)ethoxymethyl chloride (0.06ml) was then added and the mixture was warmed to room temp. over 1h. The reaction was quenched using 2M aqueous ammonia (1ml) and methanol (1ml) the solvent was evaporated and the residue was purified on a Varian Bond-Elut SPE cartridge, eluting with cyclohexane:dichloromethane (100:0 to 0:100) to give the title compound (0.67g).

20

LCMS: Rt 4.09min.

25

Intermediate 71: N-Cyclopropyl-3-[3-[(4-fluorophenyl)sulfonyl]-1-([2-(trimethylsilyl)ethyl]oxy)methyl]-1H-indazol-6-yl]-4-methylbenzamide




5

A mixture of 6-bromo-3-[(4-fluorophenyl)sulfonyl]-1-([2-(trimethylsilyl)ethyl]oxy)methyl)-1H-indazole (Intermediate 70) (0.06g) *N*-cyclopropyl-4-methyl-3-(4,4,5,5-tetramethyl-1,3,2-dioxaborolan-2-yl)benzamide (Intermediate 5) (0.04g) aqueous sodium carbonate (1M, 0.66ml) and

10 tetrakis(triphenylphosphine)palladium(0) (0.015g) in 1,2-dimethoxyethane (4ml) was stirred at reflux under nitrogen for 18h. The mixture was concentrated under vacuum and the residue was purified on a Varian Bond-Elut SPE cartridge, eluting with cyclohexane:ethyl acetate (100:0 to 50:50) to give the title compound (0.05g).

15 LCMS: Rt 3.91min.

Intermediate 72: (4-Bromo-2-nitrophenyl)methyl methyl sulfide



20


A suspension of 4-bromo-1-(bromomethyl)-2-nitrobenzene (1.0g) and sodium methanethiolate (0.285g) in tetrahydrofuran (20ml) was stirred under nitrogen for 60h. The solvent was evaporated, chloroform (50ml) was added and the residual solid was removed by filtration. The filtrate was concentrated under vacuum to give the title compound (0.89g).

25

LCMS: Rt 3.35min.

Intermediate 73: (4-Bromo-2-nitrophenyl)methyl methyl sulfone


30



A suspension of oxone (7.94g) in water (16ml) was added to a solution of (4-bromo-2-nitrophenyl)methyl methyl sulfide (Intermediate 72) (0.52g) in acetonitrile (20ml).
5 The mixture was stirred for 5h filtered and the residue was washed with ethyl acetate. The biphasic filtrate was separated, the aqueous layer was re-extracted with ethyl acetate (2 x 50ml) and the combined organic extracts were washed with aqueous sodium metabisulphite (5%, 100ml) water and brine. The solution was dried using a hydrophobic filter tube then concentrated under vacuum to give the title compound
10 (0.35g).

LCMS: Rt 2.72min.

Intermediate 74: {5-Bromo-2-[(methylsulfonyl)methyl]phenyl}amine




15

Tin(II) chloride dihydrate (1.34g) was added to a stirred suspension of (4-bromo-2-nitrophenyl)methyl methyl sulfone (Intermediate 73) in dry ethanol (10ml) and the mixture was stirred at 50° under nitrogen for 2.5h. The solvent was evaporated and the
20 residue was partitioned between aqueous sodium hydroxide (2M, 30ml) and dichloromethane (200ml). The organic phase was dried using a hydrophobic filter tube and concentrated under vacuum to give the title compound (0.26g).

LCMS: Rt 2.42min.

25


Intermediate 75: 6-Bromo-3-(methylsulfonyl)-1H-indazole



To a suspension of {5-bromo-2-[(methylsulfonyl)methyl]phenyl}amine (Intermediate 74) (0.26g) in chloroform (8ml) was added potassium acetate (0.11g) and acetic anhydride (0.195ml). The mixture was heated under nitrogen at reflux for 18h. 18-Crown-6 (0.26g) and tert-butylnitrite (0.35ml) were added and the mixture was stirred at reflux for a further 88h. The mixture was concentrated under vacuum and the residue was purified by column chromatography on silica eluting with cyclohexane:ethylacetate (4:1 to 1:1) to give the title compound (0.14g).

LCMS: Rt 2.85min.


Intermediate 76: 6-Bromo-3-(methylsulfonyl)-1-([2-(trimethylsilyl)ethoxy]methyl)-1H-indazole



Sodium tert-butoxide (0.057g) was added to an ice-cold solution of 6-bromo-3-(methylsulfonyl)-1H-indazole (Intermediate 75) (0.14g) in tetrahydrofuran (1.5ml). 2-(Trimethylsilyl)ethoxymethyl chloride (0.11ml) was then added and the mixture was warmed to room temp. over 1h. The reaction was quenched using 2M aqueous ammonia (1ml) and methanol (1ml) the solvent was evaporated and the aqueous residue was extracted with ethyl acetate. The organic extracts were dried using a hydrophobic filter tube and concentrated under vacuum to give the title compound (0.67g).

LCMS: Rt 3.81min.

Intermediate 77: N-Cyclopropyl-3-fluoro-4-methyl-5-[3-(methylsulfonyl)-1-([2-(trimethylsilyl)ethoxy]methyl)-1H-indazol-6-yl]benzamide




A mixture of 6-bromo-3-(methylsulfonyl)-1-([2-(trimethylsilyl)ethoxy]methyl)-1H-indazole (Intermediate 76) (0.18g) N-cyclopropyl-3-fluoro-4-methyl-5-(4,4,5,5-tetramethyl-1,3,2-dioxaborolan-2-yl)benzamide (Intermediate 62) (0.14g) aqueous sodium carbonate (1M, 1.32ml) and tetrakis(triphenylphosphine)palladium(0) (0.01g)

in isopropanol (2ml) was stirred at reflux under nitrogen for 18h. The mixture was concentrated under vacuum and the residue was purified by column chromatography on silica eluting with cyclohexane:ethyl acetate (2:1) to give the title compound (0.15g).

5

LCMS: Rt 3.82min.

Intermediate 78: 1-Acetyl-5-bromo-3-methyl-1H-indazole



10

A solution of 4-bromo-2-ethylaniline (1.09g) in chloroform (20ml) was stirred at room temp. then treated with potassium acetate (0.49g) followed by acetic anhydride (0.95ml). 18-Crown-6 (0.26g) and *t*-butylnitrite (1.32ml) were added after 30 min and the mixture was stirred at reflux for 18h. The reaction mixture was diluted with chloroform and washed with saturated aqueous sodium hydrogen carbonate. The organic phase was separated using a hydrophobic frit and the solvent was evaporated to give the title compound as a brown solid (1.0g).

20 NMR: δ H[CDCl₃] 8.30 (1H, d), 7.79 (1H, s), 7.63 (1H, d), 2.74 (3H, s), (2.55 (3H, s).

General Method A

The halide (30 μ moles) was dissolved in DME or DMF (0.4 - 0.5ml). The boronate ester (30 μ moles in 0.2ml DME or DMF), sodium carbonate (10% aqueous solution, 0.25ml) and

25 (A) tetrakis(triphenylphosphine) palladium (3.5 μ moles in 0.1ml DME) or

(B) FibreCatTM 1001 (12mg)

were added and the mixture was heated under nitrogen at 80° for 18h. The reaction mixture was

(A) filtered through silica (100mg) and washed with methanol

30 (B) filtered through Celite and washed with DME or DMF.

The solvent was evaporated and the residue was dissolved in DMSO (0.25ml) and purified by preparative mass-directed LC-MS.

General Method B

35

A solution of N-cyclopropyl-3-(1H-indazol-5-yl)-4-methylbenzamide (Example 17, 30mg) in dichloromethane (2ml) was added to the sulfonyl chloride (0.11mmol).

Pyridine (50 μ mol) was added and the solution was stirred at 20° for 18h (if a solution is not generated a substantial quantity of the 2-substituted product is formed). The solvent was evaporated and the residue was purified by column chromatography on silica eluting with cyclohexane:ethyl acetate (4:1 to 1:1).

5

General Method C

A solution of N-cyclopropyl-3-(1H-indazol-5-yl)-4-methylbenzamide (Example 17, or N-cyclopropyl-3-fluoro-5-(1H-indazol-5-yl)-4-methylbenzamide (Example 60) (100mg) in DMF (3ml) was treated with sodium hydride (60% dispersion in oil, 16.5mg) under nitrogen. After 10 min the sulphonyl chloride (1.2equiv) was added and the suspension was stirred at room temp. for 16h. Methanol (5ml) was added, the solvent was evaporated and the crude product was purified by column chromatography on silica eluting with a cyclohexane:ethyl acetate gradient or by reverse phase preparative Hplc.


15

General Method D

The solution of the ester methyl (5-{5-[(cyclopropylamino)carbonyl]-2-methylphenyl}-1H-indazol-1-yl)acetate (Example 117) or methyl (5-{5-[(cyclopropylamino)carbonyl]-3-fluoro-2-methylphenyl}-1H-indazol-1-yl)acetate (Example 123) (0.015g) in methanol (0.2ml) was treated with an amine (3 equiv) then stirred at 65° under nitrogen for 18h. The methanol was evaporated and the residue, in chloroform, was applied to a Varian SCX-2 ion exchange column and eluted with chloroform:methanol (9:1). If required the product was further purified by reverse phase preparative Hplc.

25

Example 1: N-Cyclopropyl-4-methyl-3-(3-piperidin-4-yl-1,2-benzisoxazol-6-yl)benzamide



Example 1 was prepared by General Method A using 6-bromo-3-piperidin-4-yl-1,2-benzisoxazole and Intermediate 5.


30

NMR: δ H [d₆-DMSO] 8.43 (1H, s), 8.08 (1H, d), 7.75 (3H, m), 7.43(2H, t), 3.6 (1H, m), 3.4(2H, m) 3.12 (2H, m), 2.85 (2H, m), 2.3 (3H, s), 2.2 (2H, m), 2.1 (2H, m), 0.67 (2H, m), 0.56 (2H, m).

LCMS: Rt 2.31min, MH⁺376.

Example 2: 4-Methyl-N-(3-morpholin-4-ylphenyl)-3-(3-piperidin-4-yl-1,2-benzisoxazol-6-yl)benzamide

5



Example 2 was prepared by General Method A using 6-bromo-3-piperidin-4-yl-1,2-benzisoxazole and Intermediate 10.


10

NMR: δ H [d₆-DMSO] 10.09 (1H, s), 8.09 (1H, d), 7.94 (1H, dd), 7.91 (1H, brs), 7.82 (1H, brs), 7.51 (1H, d), 7.47 (1H, d), 7.39 (1H, brs), 7.29 (1H, brd), 7.19 (1H, t), 6.70 (1H, dd), 3.74 (4H, apparent t), 3.48 (1H, brm), 3.29 (2H, brd), 3.09 (4H, apparent t), 2.95 (2H, brt), 2.34 (3H, s), 2.15 (2H, brd), 2.01 (2H, brt).

15

LCMS: Rt 2.56min, MH⁺497.

Example 3: N-[4-Methyl-3-(3-piperidin-4-yl-1,2-benzisoxazol-6-yl)phenyl]-2-pyrrolidin-1-ylisonicotinamide



20


Example 3 was prepared by General Method A using 6-bromo-3-piperidin-4-yl-1,2-benzisoxazole and Intermediate 13.

25

NMR: δ H [d₆-DMSO] 10.3 (1H, s), 8.4 (1H, s), 8.2 (1H, d), 8.05 (1H, d), 7.75 (3H, m), 7.4 (1H, d), 7.3 (1H, d), 6.95 (1H, d), 6.85 (1H, s), 3.45 (4H, m), 3.2 (2H, d), 2.9 (2H, t), 2.25 (3H, s), 2.1 (2H, d), 1.95 (4H, m).

LCMS: Rt 2.26min, MH⁺482.

Example 4: N-[4-Methyl-3-(3-methyl-1,2-benzisoxazol-6-yl)phenyl]-2-pyrrolidin-1-ylisonicotinamide




- 5 A mixture of 6-bromo-3-methyl-1,2-benzisoxazole (11mg), N-[4-methyl-3-(4,4,5,5-tetramethyl-[1,3,2]dioxaborolan-2-yl)-phenyl]-2-pyrrolidin-1-yl-isonicotinamide (Intermediate 13, 15mg) and aqueous sodium carbonate (0.5M, 0.25ml) in anhydrous DME (1ml) was degassed then stirred under nitrogen. A solution of
- 10 tetrakis(triphenylphosphine)palladium(0) (2.5mg) in DME (0.25ml) was added and the mixture was stirred at 80° for 16h. The solvent was removed under vacuum and the residue was partitioned between dichloromethane and water. The organic layer was separated using a hydrophobic filter, the solvent was evaporated and the residue was purified by column chromatography on silica, eluting with a cyclohexane-ethyl acetate gradient to give the title compound as an oil (5.6mg).

15 NMR δ H [d6-DMSO, 600MHz] 10.3 (1H, br.s), 8.18 (1H, d), 7.93 (1H, d), 7.74 (1H, dd), 7.71 (1H, d), 7.67 (1H, s), 7.37 (1H, d), 7.37 (1H, d), 6.97 (1H, d), 6.86 (1H, s), 3.43 (4H, br.t), 2.60 (3H, s), 2.22 (3H, s), 1.96 (4H, br.t).

LC-MS: Rt 2.91min, MH⁺ 413.

20

Example 5: N-[4-Methyl-3-(3-methyl-1,2-benzisoxazol-6-yl)phenyl]thiophene-3-carboxamide



Example 5 was prepared in a similar manner to Example 4 using 6-bromo-3-methyl-1,2-benzisoxazole (11mg) and N-[4-methyl-3-(4,4,5,5-tetramethyl-[1,3,2]dioxaborolan-2-yl)-phenyl]-3-thiopheneamide (Intermediate 15, 17mg) to give an oil (8.5mg).


5

NMR, δ H [d6-DMSO, 600MHz] 10.06 (1H, br.s), 8.32 (1H, dd), 7.93 (1H, d), 7.73 (1H, dd), 7.69 (1H, d), 7.67 (1H, br.s) 7.64 (1H, dd), 7.62 (1H, dd), 7.37 (1H, dd), 7.31 (1H, d), 2.60 (3H, s), 2.21 (3H, s).

LC-MS: Rt 3.51min, MH+ 349.

10

Example 6: N-[4-Methyl-3-(3-methyl-1,2-benzisoxazol-6-yl)phenyl]-3-furamide




Example 6 was prepared in a similar manner to Example 4 using 6-bromo-3-methyl-1,2-benzisoxazole (11mg) and N-[4-methyl-3-(4,4,5,5-tetramethyl-[1,3,2]dioxaborolan-2-yl)-phenyl]-3-furamide (Intermediate 17, 16mg) to give the title compound as an oil (8.6mg).

15

LC-MS: Rt 3.39, MH+ 333.

20

Example 7: 4-Methyl-3-(3-methyl-1,2-benzisoxazol-6-yl)-N-(3-morpholin-4-ylphenyl)benzamide




Example 7 was prepared in a similar manner to Example 4 using 6-bromo-3-methyl-1,2-benzisoxazole (11mg) and 4-methyl-N-(3-morpholin-4-yl-phenyl)-3-(4,4,5,5-tetramethyl-[1,3,2]dioxaborolan-2-yl)-benzamide (Intermediate 9, 21mg) to give the title compound as an oil (9.6mg).

NMR δ H [d₆-DMSO, 600MHz] 10.07 (1H, br.s), 7.96-7.88 (3H, m), 7.76 (1H, S), 7.50 (1H, d), 7.46 (1H, d), 7.38 (1H, br.s), 7.28 (1H, br.d), 7.18 (1H, t), 6.70 (1H, dd), 3.74 (4H, m), 2.60 (3H, s), 2.32 (3H, s).

LC-MS: Rt 3.47min, MH⁺ 428.

Example 8: 4-Methyl-3-(3-methyl-1,2-benzisoxazol-6-yl)-N-(1,3-thiazol-2-yl)benzamide




15

Example 8 was prepared in a similar manner to Example 4 using 6-bromo-3-methyl-1,2-benzisoxazole (11mg) and N-(thiazol-2-yl)-4-methyl-3-(4,4,5,5-tetramethyl-[1,3,2]dioxaborolan-2-yl)-benzamide (Intermediate 18, 17mg) followed by purification *via* mass directed autprep to give the title compound as an oil (1.3mg).

20


LC-MS: Rt 3.45min, MH⁺ 350.

Example 9: N-Cyclopropyl-4-methyl-3-(3-methyl-1,2-benzisoxazol-6-yl)benzamide



- 5 Example 9 was prepared in a similar manner to Example 4 using 6-bromo-3-methyl-1,2-benzisoxazole (11mg) and N-cyclopropyl-4-methyl-3-(4,4,5,5-tetramethyl-[1,3,2]dioxaborolan-2-yl)-benzamide (Intermediate 5, 15mg) to give the title compound as an oil (8.5mg).
- 10 NMR δ H [d₆-DMSO, 600MHz] 8.43 (1H, br.d), 7.92 (1H, d), 7.78 (1H, dd), 7.74 (1H, d), 7.70 (1H, s), 7.42-7.37 (2H, 2xd), 2.85 (1H, m), 2.60 (3H, s), 2.27 (3H, s), 0.67 (2H, m), 0.55 (2H, m).
LC-MS: Rt 3.19min, MH⁺ 307.

15 **Example 10: N-[4-Methyl-3-(3-methyl-1,2-benzisoxazol-5-yl)phenyl]-2-pyrrolidin-1-ylisonicotinamide**




- 20 Example 10 was prepared in a similar manner to Example 4 using 5-bromo-3-methyl-1,2-benzisoxazole (11mg) and N-[4-methyl-3-(4,4,5,5-tetramethyl-[1,3,2]dioxaborolan-2-yl)-phenyl]-2-pyrrolidin-1-yl-isonicotinamide (Intermediate 13, 16mg) to give the title compound as an oil (6.6).

LC-MS: Rt 2.91min, MH+ 413.

Example 11: N-Cyclopropyl-3-[3-([2-hydroxy-1-(hydroxymethyl)ethyl]amino)methyl]-1,2-benzisoxazol-6-yl]-4-methylbenzamide

5



A mixture of 2-[(6-bromo-1,2-benzisoxazol-3-yl)methyl]amino-1,3-propanediol (Intermediate 2, 8mg), N-cyclopropyl-4-methyl-3-(4,4,5,5-tetramethyl-[1,3,2]dioxaborolan-2-yl)-benzamide (Intermediate 5, 10mg) aqueous sodium carbonate (1M, 0.1ml) and tetrakis(triphenylphosphine)palladium(0) (1mg) in 2-propanol (0.2ml) was stirred at 80° under nitrogen for 18h. The solvent was evaporated and the residue was purified on a Varian Bond-Elut SPE cartridge (silica, 500mg) using a dichloromethane-methanol elution gradient (19:1 to 9:1) to give the title compound as an oil (3mg).


15

NMR δ H [d6-DMSO] 8.43 (1H, br.d), 8.08 (1H, d), 7.80 (1H, dd), 7.75 (1H, d), 7.73 (1H, br.s), 7.40 (2H, 2xd), 4.48 (2H, br.t), 4.24 (2H, s), 3.42 (4H, m), 2.85 (1H, m), 2.61 (1H, m), 2.28 (3H, s), 0.72-0.53 (4H, m).

LC-MS: Rt 2.15min, MH+ 396.

20

Example 12: N-(3-Methoxyphenyl)-4-methyl-3-(3-piperidin-4-yl-1,2-benzisoxazol-6-yl)benzamide




25

N-(3-Methoxyphenyl)-4-methyl-3-(4,4,5,5-tetramethyl-[1,3,2]dioxaborolan-2-yl)-benzamide (Intermediate 19, 54mg), DME (3ml), aqueous sodium carbonate (1M,

2ml), tetrakis (triphenylphosphine) palladium (20mg) and 6-bromo-3-piperidin-4-yl-1,2-benzisoxazole (44mg) were heated together at 80°C under nitrogen for 18h. The solvent was evaporated and the residue was purified by column chromatography on silica (10g) eluting with dichloromethane:ethanol:ammonia (40:8:1) to give the title compound (37mg).

NMR: δ H[d6-DMSO] 8.05 (1H, d), 7.88-7.97 (2H, m), 7.79 (1H, s), 7.50 (1H, d), 7.42-7.47 (2H, m), 7.36 (1H, br d), 7.23 (1H, dd), 6.67 (1H, dd), 3.09-3.14 (5H, m), 2.76 (2H, m), 2.32 (3H, s), 1.99 (2H, m), 1.83 (2H, m).
LCMS: Rt 2.66min, MH^+ 442.


Example 13: 4-Methyl-3-(3-piperidin-4-yl-1,2-benzisoxazol-6-yl)-N-(1,3,4-thiadiazol-2-yl)benzamide



4-Methyl-3-(4,4,5,5-tetramethyl-[1,3,2]dioxaborolan-2-yl)-N-(thiadiazol-2-yl)-benzamide (Intermediate 20, 50.7mg), DME (3ml), aqueous sodium carbonate (1M, 2ml), tetrakis (triphenylphosphine) palladium (20mg) and 6-bromo-3-piperidin-4-yl-1,2-benzisoxazole (44mg) were heated together at 80°C under nitrogen for 18h. The mixture was evaporated and the residue was purified by column chromatography on silica (10g) eluting with cyclohexane:ethyl acetate (8:1 to 1:1) followed by dichloromethane:ethanol:ammonia (20:8:1) to give the title compound (30.1mg).

NMR: δ H[d6-DMSO] 8.95 (1H, s), 8.01-8.09 (4H, m), 7.80 (1H, s), 7.47 (1H, d), 7.45 (1H, d), 3.25-3.47 (3H, m), 2.97 (2H, dd), 2.32 (3H, s), 2.14 (2H, m), 1.97 (2H, m).
LCMS: Rt 2.44min, MH^+ 420.

Example 14: N-[4-Methyl-3-(3-piperidin-4-yl-1,2-benzisoxazol-6-yl)phenyl]thiophene-3-carboxamide



5 N-[4-Methyl-3-(4,4,5,5-tetramethyl-[1,3,2]dioxaborolan-2-yl)-phenyl]thiophene-3-amide (Intermediate 15, 50mg), DMF (1.5ml), aqueous sodium carbonate (1M, 0.75ml), tetrakis (triphenylphosphine) palladium (7.5mg) and 6-bromo-3-piperidin-4-yl-1,2-benzisoxazole (41mg) were heated together at 80°C under nitrogen for 18h. The solvent was evaporated, and the residue was purified by column chromatography on silica (10g) eluting with dichloromethane:ethanol:ammonia (100:8:1) to give the title compound (36.8mg).


10

NMR: δ H[d6-DMSO] 10.07 (1H, d), 8.32 (1H, s), 8.00 (1H, dd), 7.59-7.74 (4H, m), 7.35 (1H, d), 7.29 (1H, dd), 6.94 (1H, d), 3.02 (2H, m), 2.66 (2H, m), 2.20 (3H, s), 1.96 (1H, m), 1.78 (2H, m), 1.50 (1H, m).

LCMS: Rt 2.71min, MH^+ 418.

15

Example 15: N-[4-Methyl-3-(3-piperidin-4-yl-1,2-benzisoxazol-6-yl)phenyl]-3-furamide



20


N-[4-Methyl-3-(4,4,5,5-tetramethyl-[1,3,2]dioxaborolan-2-yl)-phenyl]-3-furamide (Intermediate 17, 47mg), DMF (1.5ml), aqueous sodium carbonate (1M, 0.75ml), tetrakis (triphenylphosphine) palladium (7.5mg) and 6-bromo-3-piperidin-4-yl-1,2-benzisoxazole (41mg) were heated together at 80°C under nitrogen for 18h. The solvent was evaporated and the residue was purified by column chromatography on silica (10g) eluting with dichloromethane:ethanol:ammonia (100:8:1) to give the title compound (37.8mg).

25

NMR: δ H[d6-DMSO] 9.93 (1H, d), 8.35 (1H, s), 8.00 (1H, dd), 7.78 (1H, s), 7.61-7.69 (2H, m), 7.35 (1H, d), 7.28 (1H, dd), 6.98 (1H, s), 3.02 (2H, m), 2.65 (2H, m), 2.21 (3H, s), 1.96 (1H, m), 1.77 (2H, m), 1.49 (1H, m).

5 LCMS: Rt 2.55min, MH^+ 402.

Example 16: N-(Cyclopropylmethyl)-4-methyl-3-(3-piperidin-4-yl-1,2-benzisoxazol-6-yl)benzamide




10

N-Cyclopropylmethyl-4-methyl-3-(4,4,5,5-tetramethyl-[1,3,2]dioxaborolan-2-yl)-benzamide (Intermediate 21, 100mg), DME (5ml), aqueous sodium carbonate (1M, 3ml), tetrakis (triphenylphosphine) palladium (0) (37mg) and 6-bromo-3-piperidin-4-yl-1,2-benzisoxazole (76mg) were heated together at 80°C under nitrogen for 18h. The solvent was evaporated and the residue was purified by column chromatography on silica (10g) eluting with dichloromethane:ethanol:ammonia (100:8:1 to 70:8:1) to give the title compound (53mg).

20 NMR: δ H[d6-DMSO] 8.56 (1H, t), 8.03 (1H, d), 7.81 (1H, dd), 7.78 (1H, s), 7.74 (1H, s), 7.42 (1H, d), 7.38 (1H, d), 3.12 (2H, t), 3.05 (2H, m), 2.68 (2H, m), 2.29 (3H, s), 1.95 (2H, m), 1.78 (2H, m), 1.00 (1H, m), 0.41 (2H, m), 0.20 (2H, m).
LCMS: Rt 2.56min, MH^+ 390.

25 **Example 17: N-Cyclopropyl-3-(1H-indazol-5-yl)-4-methylbenzamide**



30 To a solution of tert-butyl 5-bromo-1H-indazole-1-carboxylate (Intermediate 3, 2.0g) and N-cyclopropyl-4-methyl-3-(4,4,5,5-tetramethyl-[1,3,2]dioxaborolan-2-yl)-


benzamide (Intermediate 5, 2.04g) in DME (125ml) was added tetrakis
(triphenylphosphine) palladium (0.78g) and aqueous sodium carbonate (1M, 80ml).
The mixture was heated at reflux under nitrogen for 16h. The solvent was evaporated
and the residue was purified by column chromatography on silica, eluting with
5 cyclohexane:ethyl acetate (2:1 to 1:1) to give the title compound (1.38g).

NMR: δ H [CDCl_3] 8.10 (1H, s), 7.31-7.68 (7H, m), 6.26 (1H, br s), 2.89 (1H, m),
2.29 (3H, s), 0.86 (2H, m), 0.60 (2H, m).

LCMS: Rt 2.82min, MH^+ 292.

10

Example 18: N-Cyclopropyl-4-methyl-3-[1-(phenylsulfonyl)-1H-indazol-5-yl]benzamide



15


Example 18 was prepared by General Method B using benzenesulfonyl chloride to
give the title compound (10.4mg).

NMR: δ H[d6-DMSO] 8.58 (1H, d), 8.38 (1H, d), 8.19 (1H, d), 7.97 (2H, d), 7.85 (1H,
20 s), 7.62-7.77 (6H, m), 7.39 (1H, d), 2.82 (1H, m), 2.24 (3H, s), 0.65 (2H, m), 0.53
(2H, m).

LCMS: Rt 3.37min, MH^+ 432.

25

Example 19: N-Cyclopropyl-3-[1-[(3-fluorophenyl)sulfonyl]-1H-indazol-5-yl]-4-methylbenzamide



30


Example 19 was prepared by General Method B using 3-fluorobenzenesulfonyl
chloride to give the title compound (19.2mg).

NMR: δ H[d6-DMSO] 8.62 (1H, s), 8.38 (1H, d), 8.20 (1H, d), 7.86 (1H, s), 7.60-7.84
(7H, m), 7.39 (1H, d), 2.82 (1H, m), 2.24 (3H, s), 0.65 (2H, m), 0.53 (2H, m).

LCMS: Rt 3.44min, MH^+ 450.

Example 20: 3-{1-[(3-Cyanophenyl)sulfonyl]-1H-indazol-5-yl}-N-cyclopropyl-4-methylbenzamide

5



Example 20 was prepared by General Method B using 3-cyanobenzenesulfonyl chloride to give the title compound (14.5mg).


10

NMR: $\delta H[d_6\text{-DMSO}]$ 8.64 (1H, s), 8.51 (1H, s), 8.39 (1H, d), 8.27 (1H, d), 8.21-8.24 (2H, m), 7.87 (1H, s), 7.83 (1H, dd), 7.76 (1H, d), 7.68-7.72 (2H, m), 7.39 (1H, d), 2.82 (1H, m), 2.24 (3H, s), 0.65 (2H, m), 0.51 (2H, m).

LCMS: Rt 3.34min, MH^+ 457.

15

Example 21: N-Cyclopropyl-3-{1-[(3,4-difluorophenyl)sulfonyl]-1H-indazol-5-yl}-4-methylbenzamide



20


Example 21 was prepared by General Method B using 3,4-difluorobenzenesulfonyl chloride to give the title compound (15.6mg).

NMR: $\delta H[d_6\text{-DMSO}]$ 8.63 (1H, s), 8.40 (1H, br d), 8.21 (1H, d), 8.16 (1H, td), 7.85-7.90 (2H, m), 7.65-7.80 (4H, m), 7.40 (1H, d), 2.82 (1H, m), 2.25 (3H, s), 0.67 (2H, m), 0.54 (2H, m).

25

LCMS: Rt 3.49min, MH^+ 468.

Example 22: N-Cyclopropyl-4-methyl-3-[1-(methanesulfonyl)-1H-indazol-5-yl]benzamide




5

Example 22 was prepared by General Method B using methanesulfonyl chloride to give the title compound (6.9mg).

10 NMR: δ H[d6-DMSO] 8.64 (1H, s), 8.42 (1H, d), 8.03 (1H, d), 7.91 (1H, s), 7.76 (1H, d), 7.73 (1H, s), 7.66 (1H, dd), 7.40 (1H, d), 3.50 (3H, s), 2.84 (1H, m), 2.28 (3H, s), 0.67 (2H, m), 0.55 (2H, m).

LCMS: Rt 2.97min, MH^+ 370.

15 **Example 23: N-Cyclopropyl-3-[1-(2,3-dihydro-1-benzofuran-5-ylsulfonyl)-1H-indazol-5-yl]-4-methylbenzamide**




20 Example 23 was prepared by General Method B using 2,3-dihydro-1-benzofuran-5-sulfonyl chloride to give the title compound (9.2mg).

NMR: δ H[d6-DMSO] 8.54 (1H, s), 8.39 (1H, d), 8.17 (1H, d), 7.82-7.86 (2H, m), 7.74-7.77 (2H, m), 7.70 (1H, br s), 7.66 (1H, dd), 7.39 (1H, d), 6.93 (1H, d), 4.62 (2H, dd), 3.21 (2H, dd), 2.83 (1H, m), 2.25 (3H, s), 0.66 (2H, m), 0.54 (2H, m).

25 MS: e/z + 474.

Example 24: N-Cyclopropyl-3-[1-[(3-methoxyphenyl)sulfonyl]-1H-indazol-5-yl]-4-methylbenzamide




Example 24 was prepared by General Method B using 3-methoxyphenylsulfonyl
5 chloride to give the title compound (11.0mg).

NMR: δ H[d6-DMSO] 8.59 (1H, s), 8.39 (1H, d), 8.19 (1H, d), 7.85 (1H, s), 7.76 (1H, d), 7.65-7.73 (2H, m), 7.49-7.54 (2H, m), 7.37-7.41 (2H, m), 7.30 (1H, d), 3.78 (3H, s), 2.82 (1H, m), 2.24 (3H, s), 0.66 (2H, m), 0.53 (2H, m).

10 LCMS: Rt 3.42min, MH^+ 462.

Example 25: 3-[1-(Benzylsulfonyl)-1H-indazol-5-yl]-N-cyclopropyl-4-methylbenzamide




15

Example 25 was prepared by General Method B using benzylsulfonyl chloride to give the title compound (16.4mg).

20 NMR: δ H[d6-DMSO] 8.64 (1H, s), 8.41 (1H, d), 7.80 (1H, s), 7.75 (1H, dd), 7.67 (1H, d), 7.51 (1H, d), 7.37 (1H, dd), 7.18 (1H, dd), 7.12 (2H, dd), 7.00 (2H, d), 5.02 (3H, s), 2.83 (1H, m), 2.21 (3H, s), 0.67 (2H, m), 0.54 (2H, m).

LCMS: Rt 3.34min, MH^+ 446.

25 **Example 26: N-Cyclopropyl-3-[1-[(5-isoxazol-3-ylthien-2-yl)sulfonyl]-1H-indazol-5-yl]-4-methylbenzamide**



Example 26 was prepared by General Method B using 5-isoxazol-3-ylthiophene-2-sulfonyl chloride to give the title compound (3.5mg).


5

NMR: δ H[d6-DMSO] 8.73 (1H, s), 8.70 (1H, s), 8.39 (1H, d), 8.16 (1H, d), 8.03 (1H, d), 7.90 (1H, s), 7.69-7.79 (4H, m), 7.39 (1H, d), 7.16 (1H, d), 2.83 (1H, m), 2.25 (3H, s), 0.66 (2H, m), 0.52 (2H, m).

LCMS: Rt 3.46min, MH^+ 505.

10

Example 27: 3-{1-[(4-Acetylphenyl)sulfonyl]-1H-indazol-5-yl}-N-cyclopropyl-4-methylbenzamide



15


Example 27 was prepared by General Method B using 4-acetylphenylsulfonyl chloride to give the title compound (3.6mg).

NMR: δ H[d6-DMSO] 8.62 (1H, s), 8.38 (1H, d), 8.21 (1H, d), 8.10 (4H, m), 7.86 (1H, s), 7.76 (1H, dd), 7.69-7.71 (2H, m), 7.39 (1H, d), 2.83 (1H, m), 2.57 (3H, s), 2.24 (3H, s), 0.66 (2H, m), 0.52 (2H, m).

LCMS: Rt 3.32min, MH^+ 474.

Example 28: Methyl 4-[(5-{5-[(cyclopropylamino)carbonyl]-2-methylphenyl}-1H-indazol-1-yl)sulfonyl]methyl]benzoate

25



Example 28 was prepared by General Method B using methyl 4-[(chlorosulfonyl)methyl]benzoate to give the title compound (7.8mg).


5

NMR: δ H[d6-DMSO] 8.64 (1H, s), 8.39 (1H, d), 7.78 (1H, s), 7.74 (1H, d), 7.66 (2H, d), 7.64 (1H, s), 7.45 (1H, d), 7.37 (1H, d), 7.31 (1H, d), 7.11 (2H, d), 5.17 (2H, s), 3.76 (3H, s), 2.83 (1H, m), 2.13 (3H, s), 0.67 (2H, m), 0.54 (2H, m).

LCMS: Rt 3.38min, MH^+ 504.

10

Example 29: N-Cyclopropyl-3-{1-[(3,4-dimethoxyphenyl)sulfonyl]-1H-indazol-5-yl}-4-methylbenzamide




15

Example 29 was prepared by General Method B using 3,4-dimethoxybenzenesulfonyl chloride to give the title compound (4.2mg).

20 NMR: δ H[d6-DMSO] 8.56 (1H, s), 8.39 (1H, d), 8.20 (1H, d), 7.84 (1H, s), 7.75 (1H, d), 7.70 (1H, s), 7.66 (1H, d), 7.57 (1H, dd), 7.39 (1H, d), 7.35 (1H, d), 7.14 (1H, d), 3.80 (3H, s), 3.77 (3H, s), 2.83 (1H, m), 2.24 (3H, s), 0.65 (2H, m), 0.53 (2H, m).
LCMS: Rt 3.29min, MH^+ 492.

25 **Example 30: N-Cyclopropyl-4-methyl-3-(1-{[4-(trifluoromethyl)phenyl]sulfonyl}-1H-indazol-5-yl)benzamide**



Example 30 was prepared by General Method B using
4-(trifluoromethyl)benzenesulfonyl chloride to give the title compound (13.3mg).


5

NMR: δ H[d6-DMSO] 8.64 (1H, s), 8.39 (1H, d), 8.17-8.23 (3H, m), 8.02 (2H, d), 7.87 (1H, s), 7.76 (1H, d), 7.71 (1H, d), 7.70 (1H, s), 7.39 (1H, d), 2.83 (1H, m), 2.24 (3H, s), 0.65 (2H, m), 0.53 (2H, m).

LCMS: Rt 3.61min, MH^+ 500.

10

Example 31: N-Cyclopropyl-3-{1-[(4-fluorophenyl)sulfonyl]-1H-indazol-5-yl}-4-methylbenzamide



15

Example 31 was prepared by General Method B using 4-fluorobenzenesulfonyl chloride to give the title compound (13.3mg).


NMR: δ H[d6-DMSO] 8.39 (1H, s), 8.25 (1H, d), 8.06 (2H, dd), 7.76 (1H, s), 7.71 (1H, d), 7.66 (1H, s), 7.62 (1H, d), 7.38 (1H, d), 7.29 (2H, dd), 2.82 (1H, m), 2.27 (3H, s), 0.78 (2H, m), 0.61 (2H, m).

20

LCMS: Rt 3.41min, MH^+ 450.

Example 32: 4-Methyl-3-(3-piperidin-4-yl-1,2-benzisoxazol-6-yl)-N-(1,3-thiazol-2-yl)benzamide

25




4-Methyl-3-(4,4,5,5-tetramethyl-[1,3,2]dioxaborolan-2-yl)-N-(thiazol-2-yl)-benzamide (Intermediate 18, 50mg), DME (3ml), aqueous sodium carbonate (1M, 2ml), tetrakis(triphenylphosphine)palladium (20mg) and 6-bromo-3-piperidin-4-yl-1,2-benzisoxazole (44mg, 147 μ mol) were heated together at 80°C under nitrogen for 18h. The solvent was evaporated, and the residue was purified by column chromatography on silica (10g) eluting with dichloromethane:ethanol:ammonia (100:8:1) to give the title compound (43mg).

10

NMR: δ H[d6-DMSO] 8.00-8.09 (3H, m), 7.81 (1H, s), 7.44-7.52 (3H, m), 7.19 (1H, d), 3.08 (2H, m), 2.69 (2H, m), 2.34 (3H, s), 1.98 (2H, m), 1.80 (2H, m).

LCMS: Rt 2.53min, MH^+ 419.


15 **Example 33: N-Cyclopropyl-4-methyl-3-[3-(1-piperazinyl)-1,2-benzisoxazol-6-yl]benzamide**



A solution of 1,1-dimethylethyl 4-(6-{5-[(cyclopropylamino)carbonyl]-2-methylphenyl}-1,2-benzisoxazol-3-yl)-1-piperazinecarboxylate (Intermediate 26) in methanol (1ml) was treated with a solution of hydrogen chloride in dioxan (4N, 1ml) then stirred at room temp. for 18h. The solvent was evaporated and the residue was re-dissolved in methanol and applied to an SCX cartridge (1g). Elution with methanol followed by methanol:0.88ammonia (99:1) gave the title compound as a white solid (0.016g).

25

LCMS: Rt 2.31min, MH^+ 377.

Example 34: N-Cyclopropyl-4-methyl-3-[3-(4-morpholinyl)-1,2-benzisoxazol-6-yl]benzamide

5

A mixture of 6-bromo-3-(4-morpholinyl)-1,2-benzisoxazole (Intermediate 27) (0.016g), N-Cyclopropyl-4-methyl-3-(4,4,5,5-tetramethyl-1,3,2-dioxaborolan-2-yl)benzamide (Intermediate 5), (0.03g) 2N aqueous sodium carbonate (1ml) and tetrakis(triphenylphosphine)palladium(0) (0.5mg) in isopropanol (3ml) was stirred at reflux under nitrogen for 17h. The residue was absorbed onto silica (Merck 7734) and applied to a Varian Bond-Elut SPE cartridge (silica, 5g). Elution with dichloromethane:methanol (98:2) gave the title compound as a white solid (0.024g).



10

LCMS: Rt 3.03min, MH⁺ 378.

15

Example 35: N-Cyclopropyl-4-methyl-3-[3-[2-oxo-2-(1-piperazinyl)ethyl]-1,2-benzisoxazol-6-yl]benzamide**Example 36: Methyl (6-{5-[(cyclopropylamino)carbonyl]-2-methylphenyl}-1,2-benzisoxazol-3-yl)acetate**

20



25

A suspension of 1,1-dimethylethyl 4-[(6-{5-[(cyclopropylamino)carbonyl]-2-methylphenyl}-1,2-benzisoxazol-3-yl)acetyl]-1-piperazinecarboxylate (Intermediate 34) (0.03g) in methanol (1ml) was treated with a solution of hydrogen chloride in dioxan (4M, 1ml) to give a colourless solution which was stirred at room temp. for 18h. The solvent was evaporated and the residue was partitioned between dichloromethane (2ml) and 2M aqueous sodium carbonate (2ml). The organic layer was applied to a Varian Bond-Elut SPE cartridge (silica, 5g) and sequentially eluted


with dichloromethane, ether, ethyl acetate, acetonitrile, acetone, dichloromethane:ethanol:0.88 ammonia (100:8:1) dichloromethane:ethanol:0.88 ammonia (40:8:1) ethanol and methanol to give Example 35 (6mg) as a white solid and Example 36 (4mg) as a white solid.

5

Example 35 – LCMS: Rt 2.27min, MH+ 419.

Example 36 – LCMS: Rt 3.02min, MH+ 365.

10 **Example 37: N-Cyclopropyl-3-(3-{2-[(2-hydroxyethyl)amino]-2-oxoethyl}-1,2-benzisoxazol-6-yl)-4-methylbenzamide**




15 A mixture of *N*-cyclopropyl-4-methyl-3-(4,4,5,5-tetramethyl-1,3,2-dioxaborolan-2-yl)benzamide (Intermediate 5) (0.04g) 2-(6-bromo-1,2-benzisoxazol-3-yl)-*N*-(2-hydroxyethyl)acetamide (Intermediate 35) (0.04g) 2M aqueous sodium carbonate (1.5ml) and tetrakis(triphenylphosphine)palladium(0) (1mg) in isopropanol (3ml) was stirred at reflux under nitrogen for 18h. The mixture was absorbed onto silica (Merck 7734) and applied to a Varian Bond-Elut SPE cartridge (silica, 5g) and eluted with a toluene:ethanol gradient (100:0 to 0:100) to give the title compound as a white solid (0.03g).

20

LCMS: Rt 2.60min, MH+ 394.


25 **Example 38: N-Cyclopropyl-4-methyl-3-{3-[2-oxo-2-(1-piperidinyl)ethyl]-1,2-benzisoxazol-6-yl}benzamide**



The procedure for Example 37 was followed using 6-bromo-3-[2-oxo-2-(1-piperidinyl)ethyl]-1,2-benzisoxazole (Intermediate 36) (0.04g) in place of
 5 Intermediate 35 to give the title compound as a white solid (0.03g).

LCMS: Rt 3.10min, MH+ 418.


Example 39: N-Cyclopropyl-4-methyl-3-{3-[2-(methylamino)-2-oxoethyl]-1,2-benzisoxazol-6-yl}benzamide
 10



The procedure for Example 37 was followed 2-(6-bromo-1,2-benzisoxazol-3-yl)-N-methylacetamide (Intermediate 37) (0.04g) in place of Intermediate 35 to give the title
 15 compound as a white solid (0.02g).

LCMS: Rt 2.69min, MH+ 364.


Example 40: N-Cyclopropyl-3-(3-{2-[(3-hydroxypropyl)amino]-2-oxoethyl}-1,2-benzisoxazol-6-yl)-4-methylbenzamide
 20



- 5 The procedure for Example 37 was followed 2-(6-bromo-1,2-benzisoxazol-3-yl)-N-(3-hydroxypropyl)acetamide (Intermediate 38) (0.04g) in place of Intermediate 35 to give the title compound as a white solid (0.03g).

LCMS: Rt 2.60min, MH+ 408.

10 **Example 41: N-Cyclopropyl-3-(3-{2-[(cyclopropylmethyl)amino]-2-oxoethyl}-1,2-benzisoxazol-6-yl)-4-methylbenzamide**




- 15 The procedure for Example 37 was followed 2-(6-bromo-1,2-benzisoxazol-3-yl)-N-(cyclopropylmethyl)acetamide (Intermediate 39) (0.04g) in place of Intermediate 35 to give the title compound as a white solid (0.02g).

LCMS: Rt 2.99min, MH+ 404.

20

Example 42: N-Cyclopropyl-4-methyl-3-{3-[2-oxo-2-(1-pyrrolidinyl)ethyl]-1,2-benzisoxazol-6-yl}benzamide




25

The procedure for Example 37 was followed using 6-bromo-3-[2-oxo-2-(1-pyrrolidinyl)ethyl]-1,2-benzisoxazole (Intermediate 40) (0.04g) in place of Intermediate 35 to give the title compound as a white solid (0.01g).

5 LCMS: Rt 3.19min, MH⁺ 404.


Example 43: N-Cyclopropyl-3-{3-[2-(ethylamino)-2-oxoethyl]-1,2-benzisoxazol-6-yl}-4-methylbenzamide



The procedure for Example 37 was followed using 2-(6-bromo-1,2-benzisoxazol-3-yl)-N-ethylacetamide (Intermediate 41) (0.04g) in place of Intermediate 35 to give the title compound as a white solid (0.02g).

15 LCMS: Rt 2.81min, MH⁺ 378.


20 **Example 44: N-Cyclopropyl-3-{3-[2-(cyclopropylamino)-2-oxoethyl]-1,2-benzisoxazol-6-yl}-4-methylbenzamide**



The procedure for Example 37 was followed using 2-(6-bromo-1,2-benzisoxazol-3-yl)-N-cyclopropylacetamide (Intermediate 42) (0.04g) in place of Intermediate 35 to give the title compound as a white solid (0.02g).

30 LCMS: Rt 2.83min, MH⁺ 390.

Example 45: N-Cyclopropyl-4-methyl-3-{3-[2-(4-morpholinyl)-2-oxoethyl]-1,2-benzisoxazol-6-yl}benzamide




5

The procedure for Example 37 was followed using 6-bromo-3-[2-(4-morpholinyl)-2-oxoethyl]-1,2-benzisoxazole (Intermediate 43) (0.04g) in place of Intermediate 35 to give the title compound as a white solid (0.02g).

10 LCMS: Rt 2.79min, MH⁺ 420.

Example 46: N-Cyclopropyl-4-methyl-3-{3-[2-({3-(methyloxy)phenyl}methyl)amino]-2-oxoethyl]-1,2-benzisoxazol-6-yl}benzamide



15


The procedure for Example 37 was followed using 2-(6-bromo-1,2-benzisoxazol-3-yl)-N-{{3-(methyloxy)phenyl}methyl}acetamide (Intermediate 44) (0.05g) in place of Intermediate 35 to give the title compound as a white solid (0.02g).

20

LCMS: Rt 3.16min, MH⁺ 470.

Example 47: N-Cyclopropyl-4-methyl-3-{3-[2-oxo-2-(1,3-thiazol-2-ylamino)ethyl]-1,2-benzisoxazol-6-yl}benzamide


25



- The procedure for Example 37 was followed using 2-(6-bromo-1,2-benzisoxazol-3-yl)-N-1,3-thiazol-2-ylacetamide (Intermediate 45) (0.03g) in place of Intermediate 35.
- 5 The SPE cartridge was eluted with cyclohexane:ethyl acetate (100:0 to 50:50) to give the title compound as a white solid (0.005g).

LCMS: Rt 3.06min, MH+ 433.

10 **Example 48: N-Cyclopropyl-4-methyl-3-{3-[(4-methyl-1-piperazinyl)methyl]-1,2-benzisoxazol-6-yl}benzamide**




- 15 A mixture of 6-bromo-3-[(4-methyl-1-piperazinyl)methyl]-1,2-benzisoxazole (Intermediate 46) (0.02g) *N*-cyclopropyl-4-methyl-3-(4,4,5,5-tetramethyl-1,3,2-dioxaborolan-2-yl)benzamide (Intermediate 5) (0.02g) 2N aqueous sodium carbonate and tetrakis(triphenylphosphine)palladium(0) (1mg) in isopropanol (0.3ml) was heated in a Reactivial™ at 65° for 18h. the reaction mixture was partitioned between
- 20 water and ethyl acetate, the phases were separated and the aqueous layer was re-extracted with ethyl acetate. The combined organic extracts were washed with water (x2) and brine, dried using a hydrophobic filter tube and concentrated under a stream of nitrogen. The residue was purified by reverse phase preparative Hplc to give the title compound as a white foam (0.006g).

25

LCMS: Rt 2.26min, MH+ 405.

Example 49: N-Cyclopropyl-4-methyl-3-[3-(1-piperidinylmethyl)-1,2-benzisoxazol-6-yl]benzamide




5

The procedure for Example 48 was followed using 6-bromo-3-(1-piperidinylmethyl)-1,2-benzisoxazole (Intermediate 47) (0.015g) in place of Intermediate 46 to give the title compound as a white foam (0.005g).

10 LCMS: Rt 2.28min, MH⁺ 390.

Example 50: N-Cyclopropyl-4-methyl-3-[3-(4-morpholinylmethyl)-1,2-benzisoxazol-6-yl]benzamide



15


The procedure for Example 48 was followed using 6-bromo-3-(4-morpholinylmethyl)-1,2-benzisoxazole (Intermediate 48) (0.025g) in place of Intermediate 46 to give the title compound as a white foam (0.002g).

20

LCMS: Rt 2.64min, MH⁺ 392.

Example 51: N-Cyclopropyl-4-methyl-3-[3-(1-pyrrolidinylmethyl)-1,2-benzisoxazol-6-yl]benzamide


25



- 5 The procedure for Example 48 was followed using 6-bromo-3-(1-pyrrolidinylmethyl)-1,2-benzisoxazole (Intermediate 49) (0.015g) in place of Intermediate 46 to give the title compound as a white foam (0.002g).

LCMS: Rt 2.24min, MH⁺ 376.

10 **Example 52: 3-(3-Amino-1,2-benzisoxazol-6-yl)-N-cyclopropyl-4-methylbenzamide**




- 15 A mixture of 3-amino-6-bromo-1,2-benzisoxazole (0.04g) *N*-cyclopropyl-4-methyl-3-(4,4,5,5-tetramethyl-1,3,2-dioxaborolan-2-yl)benzamide (Intermediate 5) (0.07g) and saturated aqueous sodium hydrogen carbonate (0.25ml) was stirred under nitrogen, treated with tetrakis(triphenylphosphine)palladium(0) then heated at reflux for 6h. Ethyl acetate (2ml) and water (2ml) were added to the cooled mixture and the phases
20 were separated. The organic layer was washed with brine, dried using a hydrophobic filter tube and concentrated under a stream of nitrogen. The residue was purified by reverse phase preparative Hplc to give the title compound as a white foam (0.006g).

LCMS: Rt 2.75min, MH⁺ 308.

25

Example 53: N-Cyclopropyl-3-[3-(cyclopropylamino)-1,2-benzisoxazol-6-yl]-5-fluoro-4-methylbenzamide




A mixture of 6-bromo-N-cyclopropyl-1,2-benzisoxazol-3-amine (Intermediate 53) (0.02g) N-cyclopropyl-3-fluoro-4-methyl-5-(4,4,5,5-tetramethyl-1,3,2-dioxaborolan-2-yl)benzamide (Intermediate 62) (0.02g) saturated aqueous sodium hydrogen carbonate (0.5ml) and tetrakis(triphenylphosphine)palladium(0) in isopropanol (2ml) was heated in a microwave at 150° for 10min. The residue was absorbed onto silica (Merck 7734) and applied to a Varian Bond Elut SPE cartridge (silica, 5g) eluted with cyclohexane:ethyl acetate (75:25 to 0:100) to give the title compound (10mg).

10

LCMS: Rt 3.20, MH+ 366.

Example 54: 6-{5-[(Cyclopropylamino)carbonyl]-3-fluoro-2-methylphenyl}-N-(cyclopropylmethyl)-1,2-benzisoxazole-3-carboxamide


15



A mixture of 6-bromo-N-(cyclopropylmethyl)-1,2-benzisoxazole-3-carboxamide (Intermediate 56) (0.03g) N-cyclopropyl-3-fluoro-4-methyl-5-(4,4,5,5-tetramethyl-1,3,2-dioxaborolan-2-yl)benzamide (Intermediate 62) (0.03g) saturated aqueous sodium hydrogen carbonate (0.25ml) and tetrakis(triphenylphosphine)palladium(0) (1mg) in isopropanol (0.5ml) was stirred at reflux under nitrogen for 4h. Ethyl acetate (2ml) and water (2ml) were added, the phases were separated and the aqueous layer was re-extracted with ethyl acetate. The combined organic extracts were washed with brine, dried using a hydrophobic filter tube and concentrated under a stream of nitrogen. The residue was purified on Varian Bond-Elut SPE cartridge (silica, 1g) eluting with cyclohexane:ethyl acetate (100:0 to 50:50) to give the title compound as a colourless foam (0.03g).

LCMS: Rt 3.37min, MH+ 408.


Example 55: 6-{5-[(Cyclopropylamino)carbonyl]-3-fluoro-2-methylphenyl}-N-propyl-1,2-benzisoxazole-3-carboxamide



The procedure for Example 54 was followed using 6-bromo-N-propyl-1,2-benzisoxazole-3-carboxamide (Intermediate 57) (0.04g) in place of Intermediate 56 to give the title compound as a yellow foam (0.02g).

LCMS: Rt 3.32min, MH⁺ 396.


Example 56: 6-{5-[(Cyclopropylamino)carbonyl]-3-fluoro-2-methylphenyl}-N-methyl-1,2-benzisoxazole-3-carboxamide



The procedure for Example 54 was followed using 6-bromo-N-methyl-1,2-benzisoxazole-3-carboxamide (Intermediate 58) (0.04g) in place of Intermediate 56 to give the title compound as a yellow foam (0.02g).

LCMS: Rt 3.04min, MH⁺ 368.

Example 57: 6-{5-[(Cyclopropylamino)carbonyl]-3-fluoro-2-methylphenyl}-N,N-dimethyl-1,2-benzisoxazole-3-carboxamide




The procedure for Example 54 was followed using 6-bromo-N,N-dimethyl-1,2-benzisoxazole-3-carboxamide (Intermediate 59) (0.04g) in place of Intermediate 56 to give the title compound as a yellow foam (0.02g).

LCMS: Rt 3.05min, MH+ 382.

5

Example 58: N-Cyclopropyl-6-{5-[(cyclopropylamino)carbonyl]-3-fluoro-2-methylphenyl}-1,2-benzisoxazole-3-carboxamide




10

The procedure for Example 54 was followed using 6-bromo-N-cyclopropyl-1,2-benzisoxazole-3-carboxamide (Intermediate 60) (0.04g) in place of Intermediate 56 to give the title compound as a yellow foam (0.02g).

LCMS: Rt 3.18min, MH+ 394.

15

Example 59: 1,1-Dimethylethyl 5-{5-[(cyclopropylamino)carbonyl]-3-fluoro-2-methylphenyl}-1H-indazole-1-carboxylate



20


A mixture of tert-butyl 5-bromo-1H-indazole-1-carboxylate (Intermediate 3) (1.07g) {5-[(cyclopropylamino)carbonyl]-3-fluoro-2-methylphenyl}boronic acid (Intermediate 64) (0.85g) sodium carbonate (1.9g) and tetrakis(triphenylphosphine)palladium(0) (0.42g) in 1,2-dimethoxyethane (70ml) was stirred at reflux under nitrogen for 20h. the solvent was removed and the residue was partitioned between water (50ml) and ethyl acetate (50ml). The aqueous layer was re-extracted with ethyl acetate (3x30ml) and the combined organic extracts were dried using a hydrophobic filter tube and concentrated under vacuum. The residue was purified by column chromatography on silica eluting with cyclohexane:ethyl acetate (75:25 to 60:40) to give the title compound (3.1g).

30

LCMS: Rt 3.46min, MH+ 410.

Example 60: N-Cyclopropyl-3-fluoro-5-(1H-indazol-5-yl)-4-methylbenzamide

5




A mixture of 1,1-dimethylethyl 5-{5-[(cyclopropylamino)carbonyl]-3-fluoro-2-methylphenyl}-1H-indazole-1-carboxylate (Example 59) (0.46g) in a solution of
10 hydrogen chloride in dioxan (4M, 7ml) was stirred at room temp. under nitrogen for 4.5h. The solvent was evaporated and the residue was partitioned between dichloromethane (20ml) and aqueous sodium hydroxide (2M, 20ml). The organic layer was separated using a hydrophobic filter tube, the solvent was evaporated and the residue was purified on a Varian Bond-Elut SPE cartridge (silica, 10g) eluting
15 with chloroform:methanol (100:0 to 98:2) to give the title compound (0.06g).

LCMS: Rt 2.96min, MH+ 310.

Example 61: N-Cyclopropyl-4-methyl-3-{1-[(1-methylethyl)sulfonyl]-1H-indazol-5-yl}benzamide


20



Prepared by General Method C. See Table.


25

Example 62: N-Cyclopropyl-3-[1-(ethylsulfonyl)-1H-indazol-5-yl]-4-methylbenzamide



Prepared by General Method C. See Table.


5 **Example 63: N-Cyclopropyl-3-[1-(cyclopropylsulfonyl)-1H-indazol-5-yl]-4-methylbenzamide**



Prepared by General Method C. See Table.

10


Example 64: Methyl 5-[5-[5-[(cyclopropylamino)carbonyl]-2-methylphenyl]-1H-indazol-1-yl)sulfonyl]-2-furancarboxylate



15

Prepared by General Method C. See Table.

Example 65: 3-[1-(2,1,3-Benzoxadiazol-4-ylsulfonyl)-1H-indazol-5-yl]-N-cyclopropyl-4-methylbenzamide




5

Prepared by General Method C. See Table.


Example 66: N-Cyclopropyl-3-{1-[(2-fluorophenyl)sulfonyl]-1H-indazol-5-yl}-4-methylbenzamide

10




Prepared by General Method C. See Table.

Example 67: N-Cyclopropyl-4-methyl-3-{1-[(4-methylphenyl)sulfonyl]-1H-indazol-5-yl}benzamide




Prepared by General Method C. See Table.

5 **Example 68: N-Cyclopropyl-4-methyl-3-[1-(2-thienylsulfonyl)-1H-indazol-5-yl]benzamide**



10 Prepared by General Method C. See Table.


Example 69: N-Cyclopropyl-3-[1-[(3,5-dimethyl-4-isoxazolyl)sulfonyl]-1H-indazol-5-yl]-4-methylbenzamide



15


Prepared by General Method C. See Table.

Example 70: N-Cyclopropyl-4-methyl-3-[1-(propylsulfonyl)-1H-indazol-5-yl]benzamide




Prepared by General Method C. See Table.

10 **Example 71: N-Cyclopropyl-4-methyl-3-{1-[(2-nitrophenyl)sulfonyl]-1H-indazol-5-yl}benzamide**



15 Prepared by General Method C. See Table.


Example 72: N-Cyclopropyl-4-methyl-3-{1-[(2,2,2-trifluoroethyl)sulfonyl]-1H-indazol-5-yl}benzamide



Prepared by General Method C. See Table.

Example 73: 3-{1-[(2-Cyanophenyl)sulfonyl]-1H-indazol-5-yl}-N-cyclopropyl-4-methylbenzamide


5



Prepared by General Method C. See Table.

Example 74: N-Cyclopropyl-4-methyl-3-{1-[(1,3,5-trimethyl-1H-pyrazol-4-yl)sulfonyl]-1H-indazol-5-yl}benzamide


10



Prepared by General Method C. See Table.


Example 75: N-Cyclopropyl-4-methyl-3-{1-[(1-methyl-1H-imidazol-4-yl)sulfonyl]-1H-indazol-5-yl}benzamide

15



Prepared by General Method C. See Table.


5 **Example 76: N-Cyclopropyl-3-fluoro-4-methyl-5-[1-(2-thienylsulfonyl)-1H-indazol-5-yl]benzamide**



Prepared by General Method C. See Table.

10

Example 77: N-Cyclopropyl-3-fluoro-4-methyl-5-[1-(3-thienylsulfonyl)-1H-indazol-5-yl]benzamide




15

Prepared by General Method C. See Table.


Example 78: N-Cyclopropyl-3-fluoro-4-methyl-5-[1-[(4-methylphenyl)sulfonyl]-1H-indazol-5-yl]benzamide

20



Prepared by General Method C. See Table.

Example 79: N-Cyclopropyl-3-fluoro-4-methyl-5-(1-{[4-(methoxy)phenyl]sulfonyl}-1H-indazol-5-yl)benzamide




5

Prepared by General Method C. See Table.

Example 80: N-Cyclopropyl-3-fluoro-4-methyl-5-(1-{[4-(1-methylethyl)phenyl]sulfonyl}-1H-indazol-5-yl)benzamide


10



Prepared by General Method C. See Table.


Example 81: 3-{1-[4-Chlorophenyl]sulfonyl}-1H-indazol-5-yl}-N-cyclopropyl-5-fluoro-4-methylbenzamide

15




20 Prepared by General Method C. See Table.

Example 82: N-Cyclopropyl-3-{1-[2,4-difluorophenyl]sulfonyl}-1H-indazol-5-yl}-5-fluoro-4-methylbenzamide




Prepared by General Method C. See Table.

5 **Example 83: 3-{1-[(3-Chlorophenyl)sulfonyl]-1H-indazol-5-yl}-N-cyclopropyl-5-fluoro-4-methylbenzamide**



10 Prepared by General Method C. See Table.


Example 84: N-Cyclopropyl-3-fluoro-5-{1-[(5-fluoro-2-methylphenyl)sulfonyl]-1H-indazol-5-yl}-4-methylbenzamide



15


Prepared by General Method C. See Table.

20 **Example 85: N-Cyclopropyl-3-fluoro-5-{1-[(3-fluorophenyl)sulfonyl]-1H-indazol-5-yl}-4-methylbenzamide**




Prepared by General Method C. See Table.

5 **Example 86: N-Cyclopropyl-3-fluoro-4-methyl-5-(1-([3-(methyloxy)phenyl]sulfonyl)-1H-indazol-5-yl)benzamide**



10 Prepared by General Method C. See Table.


Example 87: N-Cyclopropyl-3-[1-[(3,5-difluorophenyl)sulfonyl]-1H-indazol-5-yl]-5-fluoro-4-methylbenzamide



15


Prepared by General Method C. See Table.

20 **Example 88: N-Cyclopropyl-3-fluoro-4-methyl-5-(1-[(3-methylphenyl)sulfonyl]-1H-indazol-5-yl)benzamide**




Prepared by General Method C. See Table.

5 **Example 89: N-Cyclopropyl-3-fluoro-4-methyl-5-[1-[(1-methylethyl)sulfonyl]-1H-indazol-5-yl]benzamide**



10 Prepared by General Method C. See Table.


Example 90: N-Cyclopropyl-3-fluoro-4-methyl-5-[1-(propylsulfonyl)-1H-indazol-5-yl]benzamide



15


Prepared by General Method C. See Table.

20 **Example 91: N-Cyclopropyl-3-fluoro-4-methyl-5-[1-(methylsulfonyl)-1H-indazol-5-yl]benzamide**



Prepared by General Method C. See Table.


5 **Example 92: 3-[1-(Butylsulfonyl)-1H-indazol-5-yl]-N-cyclopropyl-5-fluoro-4-methylbenzamide**



Prepared by General Method C. See Table.

10


Example 93: N-Cyclopropyl-3-fluoro-4-methyl-5-[1-(octylsulfonyl)-1H-indazol-5-yl]benzamide



15


Prepared by General Method C. See Table.

Example 94: N-Cyclopropyl-3-[1-(cyclopropylsulfonyl)-1H-indazol-5-yl]-5-fluoro-4-methylbenzamide




Prepared by General Method C. See Table.

5 **Example 95: 3-{1-[(5-Chloro-2-thienyl)sulfonyl]-1H-indazol-5-yl}-N-cyclopropyl-5-fluoro-4-methylbenzamide**



10 Prepared by General Method C. See Table.


Example 96: N-Cyclopropyl-3-{1-[(3,5-dimethyl-4-isoxazolyl)sulfonyl]-1H-indazol-5-yl}-5-fluoro-4-methylbenzamide



15


Prepared by General Method C. See Table.

20 **Example 97: N-Cyclopropyl-3-{1-[(1,2-dimethyl-1H-imidazol-4-yl)sulfonyl]-1H-indazol-5-yl}-5-fluoro-4-methylbenzamide**




Prepared by General Method C. See Table.

5 **Example 98: N-Cyclopropyl-3-[1-(ethylsulfonyl)-1H-indazol-5-yl]-5-fluoro-4-methylbenzamide**



10 Prepared by General Method C. See Table.


Example 99: 3-{1-[2-Chlorophenyl)sulfonyl]-1H-indazol-5-yl}-N-cyclopropyl-5-fluoro-4-methylbenzamide



15


Prepared by General Method C. See Table.

20 **Example 100: N-Cyclopropyl-3-fluoro-4-methyl-5-{1-[2-methylphenyl)sulfonyl]-1H-indazol-5-yl}benzamide**




Prepared by General Method C. See Table.

5 **Example 101: N-Cyclopropyl-3-{1-[(4-ethylphenyl)sulfonyl]-1H-indazol-5-yl}-5-fluoro-4-methylbenzamide**



10 Prepared by General Method C. See Table.


Example 102: N-Cyclopropyl-3-fluoro-4-methyl-5-{1-[(4-propylphenyl)sulfonyl]-1H-indazol-5-yl}benzamide



15


Prepared by General Method C. See Table.

20 **Example 103: N-Cyclopropyl-3-{1-[(3,4-difluorophenyl)sulfonyl]-1H-indazol-5-yl}-5-fluoro-4-methylbenzamide**




Prepared by General Method C. See Table.

5 **Example 104: 3-{1-[(4-Butylphenyl)sulfonyl]-1H-indazol-5-yl}-N-cyclopropyl-5-fluoro-4-methylbenzamide**



10 Prepared by General Method C. See Table.


Example 105: N-Cyclopropyl-3-fluoro-4-methyl-5-[1-(phenylsulfonyl)-1H-indazol-5-yl]benzamide



15


Prepared by General Method C. See Table.

20 **Example 106: N-Cyclopropyl-3-fluoro-4-methyl-5-{1-[(phenylmethyl)sulfonyl]-1H-indazol-5-yl}benzamide**




Prepared by General Method C. See Table.

5 **Example 107: N-Cyclopropyl-3-fluoro-4-methyl-5-(1-[(E)-2-phenylethenyl]sulfonyl)-1H-indazol-5-yl)benzamide**



10 Prepared by General Method C. See Table.


Example 108: N-Cyclopropyl-3-{1-[(2,5-difluorophenyl)sulfonyl]-1H-indazol-5-yl}-5-fluoro-4-methylbenzamide



15


Prepared by General Method C. See Table.

20 **Example 109: N-Cyclopropyl-3-fluoro-5-{1-[(2-fluorophenyl)sulfonyl]-1H-indazol-5-yl}-4-methylbenzamide**




Prepared by General Method C. See Table.

5 **Example 110: 3-{1-[(4-Cyanophenyl)sulfonyl]-1H-indazol-5-yl}-N-cyclopropyl-5-fluoro-4-methylbenzamide**



10 Prepared by General Method C. See Table.

Example 111: N-Cyclopropyl-3-[1-(2,3-dihydro-1-benzofuran-5-ylsulfonyl)-1H-indazol-5-yl]-5-fluoro-4-methylbenzamide



15

Prepared by General Method C. See Table.

Table (Examples 61 – 111)

20

| Example No | Indazole Precursor | Sulphonyl Chloride | Rt (min) | MH ⁺ |
|------------|-----------------------|---|-------------|-----------------|
| Example 61 | Example 17 | 2-propanesulfonyl chloride ¹ | 3.17 | 398 |

| | | | |
|-------------|--|------|-----|
| Example 62 | Example 17 ethanesulfonyl chloride ² | 3.13 | 384 |
| Example 63 | Example 17 cyclopropanesulfonyl chloride ³ | 3.18 | 396 |
| Example 64 | Example 17 methyl 5-(chlorosulfonyl)-2-furancarboxylate ⁴ | 3.29 | 480 |
| Example 65 | Example 17 2,1,3-benzoxadiazole-4-sulfonyl chloride ⁴ | 3.36 | 474 |
| Example 66 | Example 17 2-fluorobenzenesulfonyl chloride ¹ | 3.38 | 450 |
| Example 67 | Example 17 4-methylbenzenesulfonyl chloride ¹ | 3.38 | 446 |
| Example 68 | Example 17 2-thiophenesulfonyl chloride ¹ | 3.33 | 438 |
| Example 69 | Example 17 3,5-dimethyl-4-isoxazolesulfonyl chloride ⁴ | 3.38 | 451 |
| Example 70 | Example 17 1-propanesulfonyl chloride ⁵ | 3.11 | 398 |
| Example 71 | Example 17 2-nitrobenzenesulfonyl chloride ¹ | 3.4 | 477 |
| Example 72 | Example 17 2,2,2-trifluoroethanesulfonyl chloride ¹ | 3.11 | 438 |
| Example 73 | Example 17 2-cyanobenzenesulfonyl chloride ¹ | 3.31 | 457 |
| Example 74 | Example 17 1,3,5-trimethyl-1 <i>H</i> -pyrazole-4-sulfonyl chloride ⁴ | 3.18 | 464 |
| Example 75 | Example 17 1-methyl-1 <i>H</i> -imidazole-4-sulfonyl chloride ⁴ | 2.88 | 436 |
| Example 76 | Example 60 2-thiophenesulfonyl chloride ¹ | 3.42 | 456 |
| Example 77 | Example 60 3-thiophenesulfonyl chloride ⁴ | 3.39 | 456 |
| Example 78 | Example 60 4-methylbenzenesulfonyl chloride ¹ | 3.57 | 464 |
| Example 79 | Example 60 4-(methyloxy)benzenesulfonyl chloride ² | 3.5 | 480 |
| Example 80 | Example 60 4-(1-methylethyl)benzenesulfonyl chloride ¹ | 3.77 | 492 |
| Example 81 | Example 60 4-chlorobenzenesulfonyl chloride ¹ | 3.64 | 484 |
| Example 82 | Example 60 2,4-difluorobenzenesulfonyl chloride ¹ | 3.53 | 486 |
| Example 83 | Example 60 3-chlorobenzenesulfonyl chloride ¹ | 3.65 | 484 |
| Example 84 | Example 60 5-fluoro-2-methylbenzenesulfonyl chloride ¹ | 3.65 | 482 |
| Example 85 | Example 60 3-fluorobenzenesulfonyl chloride ¹ | 3.53 | 468 |
| Example 86 | Example 60 3-(methyloxy)benzenesulfonyl chloride ¹ | 3.65 | 480 |
| Example 87 | Example 60 3,5-difluorobenzenesulfonyl chloride ¹ | 3.73 | 486 |
| Example 88 | Example 60 3-methylbenzenesulfonyl chloride ² | 3.71 | 464 |
| Example 89 | Example 60 4-(methyloxy)benzenesulfonyl chloride ² | 3.35 | 416 |
| Example 90 | Example 60 1-propanesulfonyl chloride ⁵ | 3.39 | 416 |
| Example 91 | Example 60 methanesulfonyl chloride ² | 3.14 | 388 |
| Example 92 | Example 60 1-butan sulfonyl chloride ¹ | 3.46 | 430 |
| Example 93 | Example 60 1-octanesulfonyl chloride ² | 3.95 | 486 |
| Example 94 | Example 60 cyclopropanesulfonyl chloride ³ | 3.25 | 414 |
| Example 95 | Example 60 5-chloro-2-thiophenesulfonyl chloride ¹ | 3.68 | 490 |
| Example 96 | Example 60 3,5-dimethyl-4-isoxazolesulfonyl chloride ² | 3.48 | 469 |
| Example 97 | Example 60 1,2-dimethyl-1 <i>H</i> -imidazole-4-sulfonyl chloride ⁴ | 3 | 468 |
| Example 98 | Example 60 ethanesulfonyl chloride ² | 3.21 | 402 |
| Example 99 | Example 60 2-chlorobenzenesulfonyl chloride ¹ | 3.59 | 484 |
| Example 100 | Example 60 2-methylbenzenesulfonyl chloride ¹ | 3.59 | 464 |
| Example 101 | Example 60 4-ethylbenzenesulfonyl chloride ⁴ | 3.68 | 478 |
| Example 102 | Example 60 4-propylbenzenesulfonyl chloride ² | 3.81 | 492 |
| Example 103 | Example 60 3,4-difluorobenzenesulfonyl chloride ¹ | 3.59 | 486 |
| Example 104 | Example 60 4-butylbenzenesulfonyl chloride ² | 3.9 | 506 |

| | | | |
|-------------|--|------|-----|
| Example 105 | Example 60 benzenesulfonyl chloride ¹ | 3.45 | 450 |
| Example 106 | Example 60 phenylmethanesulfonyl chloride ¹ | 3.42 | 464 |
| Example 107 | Example 60 (<i>E</i>)-2-phenylethenesulfonyl chloride ¹ | 3.58 | 476 |
| Example 108 | Example 60 2,5-difluorobenzenesulfonyl chloride ¹ | 3.52 | 486 |
| Example 109 | Example 60 2-fluorobenzenesulfonyl chloride ¹ | 3.47 | 468 |
| Example 110 | Example 60 4-cyanobenzenesulfonyl chloride ² | 3.42 | 475 |
| Example 111 | Example 60 2,3-dihydro-1-benzofuran-5-sulfonyl chloride ⁴ | 3.47 | 492 |

1 Purchased from Aldrich.


2 Purchased from Avocado.

3 Purchased from Array Biopharma Inc.

5 4 Purchased from Maybridge.

5 Purchased from Pfaltz & Bauer.

Example 112: N-Cyclopropyl-3-fluoro-4-methyl-5-[1-[(4-methylphenyl)sulfonyl]-1H-indol-5-yl]benzamide



10


A mixture of 5-bromo-1-[(4-methylphenyl)sulfonyl]-1H-indole (0.035g) {5-[(cyclopropylamino)carbonyl]-3-fluoro-2-methylphenyl}boronic acid (Intermediate 64) (0.024g) saturated aqueous sodium hydrogen carbonate (0.5ml) and tetrakis(triphenylphosphine)palladium(0) (2mg) in isopropanol (2ml) was heated in a microwave at 150° for 10min. The solid was removed by filtration and the residue was purified by preparative Hplc to give the title compound (0.013g).

15

LCMS: Rt 3.71min, MH+ 463.

20


Example 113: N-Cyclopropyl-3-fluoro-4-methyl-5-[1-(phenylsulfonyl)-1H-indol-5-yl]benzamide



The procedure for Example 112 was followed using 5-bromo-1-(phenylsulfonyl)-1H-indole (0.033g) in place of 5-bromo-1-[(4-methylphenyl)sulfonyl]-1H-indole to give the title compound (0.009g).

LCMS: Rt 3.61min, MH+449.


Example 114: N-Cyclopropyl-3-[1-(cyclopropylmethyl)-1H-indazol-5-yl]-4-methylbenzamide



A solution of N-cyclopropyl-3-(1H-indazol-5-yl)-4-methylbenzamide (Example 17) (0.10g) in dry DMF (3ml) was treated with sodium hydride (60% oil dispersion, 0.017g) under nitrogen. After 5min cyclopropylmethyl bromide (0.04ml) was added and the yellow solution was stirred at room temp. for 18h. Methanol (2drops) was added and the solvent was evaporated. The residue was purified by column chromatography on silica eluting with cyclohexane:ethyl acetate (2:1 to 1:1) to give the title compound (0.05g).

LCMS: Rt 3.21min, MH+ 346.

Example 115: 1,1-Dimethylethyl 4-(5-{5-[(cyclopropylamino)carbonyl]-2-methylphenyl}-1H-indazol-1-yl)-1-piperidinecarboxylate




A solution of N-cyclopropyl-3-(1H-indazol-5-yl)-4-methylbenzamide (Example 17) (0.07g) in dry DMF (3ml) was treated with sodium hydride (60% oil dispersion, 0.014g) under nitrogen. After 5min 1,1-dimethylethyl 4-[(methylsulfonyl)oxy]-1-piperidinecarboxylate (0.08g) was added and the mixture was stirred at room temp. for 18h. 0.88 Ammonia (1ml) was added and the mixture was concentrated under vacuum. The residue was purified by column chromatography on silica eluting with cyclohexane:ethyl acetate (1:1) followed by reverse phase preparative Hplc to give the title compound (0.005g).

LCMS: Rt 3.55min, MH+ 475.

10

Example 116: N-Cyclopropyl-4-methyl-3-[1-(4-piperidinyl)-1H-indazol-5-yl]benzamide hydrochloride



15


A suspension of 1,1-dimethylethyl 4-(5-{5-[(cyclopropylamino)carbonyl]-2-methylphenyl}-1H-indazol-1-yl)-1-piperidinecarboxylate (Example 115) (0.01g) in a solution of hydrogen chloride in dioxan (4M, 1ml) was treated with methanol (0.25ml) to give a solution which was stirred at room temp. for 18h. The solvent was evaporated and the residue was dried under vacuum to give the title compound (0.009g).

20

LCMS: Rt 2.40min, MH+ 375.

25


Example 117: Methyl (5-{5-[(cyclopropylamino)carbonyl]-2-methylphenyl}-1H-indazol-1-yl)acetate



- 5 A solution of N-cyclopropyl-3-(1H-indazol-5-yl)-4-methylbenzamide (Example 17) (0.65g) in dry DMF (26ml) was treated with sodium hydride (60% oil dispersion, 0.134g) under nitrogen. After 5min methyl chloroacetate (0.29g) was added and the mixture was stirred for 2h. Aqueous ammonia (2M, 10ml) was added and the solvent was removed under vacuum. The residue was purified by column chromatography on silica eluting with cyclohexane:ethyl acetate (2:1 to 1:1) to give the title compound (0.56g).

LCMS: Rt 2.95min, MH+ 364.


10 **Example 118: N-Cyclopropyl-3-(1H-indazol-6-yl)-4-methylbenzamide**



- 15 A mixture of 1-acetyl-6-bromo-1H-indazole (Intermediate 65) (0.50g) N-cyclopropyl-4-methyl-3-(4,4,5,5-tetramethyl-1,3,2-dioxaborolan-2-yl)benzamide (Intermediate 5) (0.59g) aqueous sodium carbonate(1M, 2.35ml) and PdCl₂dppf (0.237g) in 1,2-dimethoxyethane (30ml) was stirred at reflux under nitrogen for 16h. More sodium carbonate (2.3ml) and catalyst (120mg) were added and reflux was continued for a further 4h. Conc. hydrochloric acid (1ml) was added and the mixture was heated for a further 1h. The mixture was absorbed onto silica then purified by column chromatography on silica eluting with cyclohexane:ethyl acetate (2:1 to 1:4) to give the title compound (0.25g).

LCMS: Rt 2.88min, MH+ 292.


25 **Example 119: N-Cyclopropyl-3-{3-[(4-fluorophenyl)sulfonyl]-1H-indazol-6-yl}-4-methylbenzamide**



- 5 To a solution of N-cyclopropyl-3-[3-[(4-fluorophenyl)sulfonyl]-1-([2-(trimethylsilyl)ethyl]oxy)methyl]-1H-indazol-6-yl]-4-methylbenzamide (Intermediate 71) (0.05g) in methanol (1.5ml) was added conc hydrochloric acid (0.75ml). The solution was heated at reflux for 15h then left at room temp. for 18h. The solution was concentrated under vacuum then purified on a Varian Bond-Elut SPE cartridge (silica, 5g) eluting with cyclohexane:ethyl acetate (1:1) to give the title compound (0.01g).

LCMS: Rt 3.32 min, MH+ 450.

10 **Example 120: N-Cyclopropyl-3-fluoro-4-methyl-5-[3-(methylsulfonyl)-1H-indazol-6-yl]benzamide**




- 15 Conc. hydrochloric acid ((0.5ml) was added to a solution of N-cyclopropyl-3-fluoro-4-methyl-5-[3-(methylsulfonyl)-1-([2-(trimethylsilyl)ethyl]oxy)methyl]-1H-indazol-6-yl]benzamide (Intermediate 77) (0.15g) then stirred at reflux for 12h. The solvent was evaporated and the residue was purified by column chromatography on silica, eluting with cyclohexane:ethyl acetate (1:1 to 1:2) to give the title compound (0.085g).

20

LCMS: Rt 2.91min, MH+ 388.

Example 121: N-Cyclopropyl-3-fluoro-4-methyl-5-(3-methyl-1H-indazol-5-yl)benzamide

25




A mixture of 1-acetyl-5-bromo-3-methyl-1H-indazole (0.100g) {5-[(cyclopropylamino)carbonyl]-3-fluoro-2-methylphenyl}boronic acid (Intermediate

64) (0.094g) saturated aqueous sodium hydrogen carbonate (2ml) and tetrakis(triphenylphosphine)palladium(0) (9mg) in isopropanol (6ml) was heated in a microwave at 150° for 10min. The solvent was evaporated and the residue was purified on a Varian Bond-Elut SPE cartridge (silica 10g) eluting with cyclohexane:ethyl acetate(1:1) to give the title compound (0.073g).

LCMS: Rt 3.02min, MH+ 324.


Example 122: N-Cyclopropyl-3-fluoro-4-methyl-5-[3-methyl-1-(2-thienylsulfonyl)-1H-indazol-5-yl]benzamide



Sodium hydride (60% oil dispersion, 0.008g) was added to a stirred solution of *N*-cyclopropyl-3-fluoro-4-methyl-5-(3-methyl-1*H*-indazol-5-yl)benzamide (Example 121) (0.033g) in dry DMF(1.5ml) under nitrogen. After 5min a solution of 2-thiophenesulphonyl chloride (0.038g) in DMF (0.75ml) was added and stirring was continued for a further 2h. Water (1ml) and chloroform (10ml) were added, the mixture was passed through a hydrophobic filter and the solvent was evaporated. The residue was purified by preparative Hplc to give the title compound as a clear glass (0.024g).

LCMS: Rt 3.44min, MH+ 470.

Example 123: Methyl (5-[5-[(cyclopropylamino)carbonyl]-3-fluoro-2-methylphenyl]-1H-indazol-1-yl)acetate




A solution of N-cyclopropyl-3-fluoro-5-(1H-indazol-5-yl)-4-methylbenzamide
 (Example 60) (0.35g) in dry DMF (5ml) was treated with sodium hydride (60% oil
 dispersion, 0.08g) at room temp under nitrogen then stirred for 20min. A solution of
 methyl bromoacetate (0.2ml) in dry DMF (0.5ml) was added and stirring was
 5 continued for 1h. Water (30ml) was added and the mixture was extracted with ethyl
 acetate (x2). The organic extracts were washed with water (x3) and brine, dried
 (Na₂SO₄) and concentrated under vacuum. The residue was purified using a Varian
 Bond-Elut SPE cartridge (silica, 10g) eluting with cyclohexane:ethyl acetate (4:1
 to 1:1) to give the title compound as a yellow foam (0.26g).

10

LCMS: Rt 3.03min, MH⁺ 382.

Example 124: N-Cyclopropyl-4-methyl-3-{1-[2-({[4-(methoxy)phenyl]methyl}amino)-2-oxoethyl]-1H-indazol-5-yl}benzamide


15



Example 124 was prepared by General Method D using methyl (5-{5-
 [(cyclopropylamino)carbonyl]-2-methylphenyl}-1H-indazol-1-yl)acetate (Example
 117) and 4-methoxybenzylamine to give the title compound (0.011g).

LCMS: Rt 2.99min, MH⁺ 469.


Example 125: N-Cyclopropyl-4-methyl-3-(1-{2-oxo-2-[(phenylmethyl)amino]ethyl}-1H-indazol-5-yl)benzamide



Example 125 was prepared by General Method D using methyl (5-{5-
[(cyclopropylamino)carbonyl]-2-methylphenyl}-1H-indazol-1-yl)acetate (Example
5 117) and benzylamine to give the title compound (0.002g).

LCMS: Rt 3.00min, MH+ 439.

Example 126: N-Cyclopropyl-4-methyl-3-[1-(2-{[(4-methylphenyl)methyl]amino}-2-oxoethyl)-1H-indazol-5-yl]benzamide




10 Example 126 was prepared by General Method D using methyl (5-{5-
[(cyclopropylamino)carbonyl]-2-methylphenyl}-1H-indazol-1-yl)acetate (Example
117) and 4-methylbenzylamine to give the title compound (0.010g)

LCMS: Rt 3.11min, MH+ 453.

Example 127: N-Cyclopropyl-4-methyl-3-(1-(2-{[(4-methylphenyl)methyl]amino}-2-oxoethyl)-1H-indazol-5-yl)benzamide


15



Example 127 was prepared by General Method D using methyl (5-{5-[(cyclopropylamino)carbonyl]-2-methylphenyl}-1H-indazol-1-yl)acetate (Example 117) and 4-(aminomethyl)-N-methylbenzamide to give the title compound (0.005g).

5 LCMS: Rt 2.68min, MH+ 496.


Example 128: N-Cyclopropyl-4-methyl-3-{1-[2-oxo-2-(propylamino)ethyl]-1H-indazol-5-yl}benzamide



10 Example 128 was prepared by General Method D using methyl (5-{5-[(cyclopropylamino)carbonyl]-2-methylphenyl}-1H-indazol-1-yl)acetate (Example 117) and 1-propylamine to give the title compound (0.011g).

LCMS: Rt 2.81min, MH+ 391.


Example 129: N-Cyclopropyl-3-(1-{2-[(cyclopropylmethyl)amino]-2-oxoethyl}-1H-indazol-5-yl)-4-methylbenzamide



Example 129 was prepared by General Method D using methyl (5-{5-[(cyclopropylamino)carbonyl]-2-methylphenyl}-1H-indazol-1-yl)acetate (Example 117) and 1-cyclopropylmethanamine to give the title compound (0.015g).

5 LCMS: Rt 2.86min, MH+ 403.


Example 130: N-Cyclopropyl-3-(1-(2-[(2,2-dimethylpropyl)amino]-2-oxoethyl)-1H-indazol-5-yl)-4-methylbenzamide



10 Example 130 was prepared by General Method D using methyl (5-{5-[(cyclopropylamino)carbonyl]-2-methylphenyl}-1H-indazol-1-yl)acetate (Example 117) and 2,2-dimethyl-1-propanamine to give the title compound (0.015g).

LCMS: Rt 3.06min, MH+ 419.


Example 131: N-Cyclopropyl-4-methyl-3-{1-[2-(4-morpholinyl)-2-oxoethyl]-1H-indazol-5-yl}benzamide



Example 131 was prepared by General Method D using methyl (5-{5-
5 [(cyclopropylamino)carbonyl]-2-methylphenyl}-1H-indazol-1-yl)acetate (Example
117) and morpholine to give the title compound (0.002g).

LCMS: Rt 2.66min, MH+ 419.


10 **Example 132: N-Cyclopropyl-3-fluoro-4-methyl-5-(1-{2-oxo-2-
[(phenylmethyl)amino]ethyl}-1H-indazol-5-yl)benzamide**



Example 132 was prepared by General Method D using methyl (5-{5-
15 [(cyclopropylamino)carbonyl]-3-fluoro-2-methylphenyl}-1H-indazol-1-yl)acetate
(Example 123) and benzylamine to give the title compound (0.008g).

LCMS: Rt 3.17min, MH+ 457.


Example 133: N-Cyclopropyl-3-fluoro-4-methyl-5-[1-(2-[[4-methylphenyl)methyl]amino]-2-oxoethyl]-1H-indazol-5-yl]benzamide



- Example 133 was prepared by General Method D using methyl (5-{5-
5 [(cyclopropylamino)carbonyl]-3-fluoro-2-methylphenyl}-1H-indazol-1-yl)acetate
(Example 123) and 4-methylbenzylamine to give the title compound (0.011g).

LCMS: Rt 3.28min, MH⁺ 471.


- 10 **Example 134: N-Cyclopropyl-3-fluoro-4-methyl-5-[1-(2-[[4-methoxyphenyl)methyl]amino]-2-oxoethyl]-1H-indazol-5-yl]benzamide**



- Example 134 was prepared by General Method D using methyl (5-{5-
15 [(cyclopropylamino)carbonyl]-3-fluoro-2-methylphenyl}-1H-indazol-1-yl)acetate
(Example 123) and 4-methoxybenzylamine to give the title compound (0.012g).

LCMS: Rt 3.15min, MH⁺ 487.

Example 135: N-Cyclopropyl-3-fluoro-4-methyl-5-(1-{2-[(4-[(methylamino)carbonyl]phenyl)methyl]amino}-2-oxoethyl)-1H-indazol-5-yl}benzamide




5

Example 135 was prepared by General Method D using using methyl (5-{5-[(cyclopropylamino)carbonyl]-3-fluoro-2-methylphenyl}-1*H*-indazol-1-yl)acetate (Example 123) and 4-(aminomethyl)-*N*-methylbenzamide to give the title compound (0.005g).

10 LCMS: Rt 2.80min, MH+ 514.


Example 136: N-Cyclopropyl-3-fluoro-4-methyl-5-{1-[2-oxo-2-(propylamino)ethyl]-1H-indazol-5-yl}benzamide



15 Example 136 was prepared by General Method D using using methyl (5-{5-[(cyclopropylamino)carbonyl]-3-fluoro-2-methylphenyl}-1*H*-indazol-1-yl)acetate (Example 123) and 1-propylamine to give the title compound (0.005g).

LCMS: Rt 2.92min, MH+ 409.


Example 137: N-Cyclopropyl-3-(1-{2-[(cyclopropylmethyl)amino]-2-oxoethyl}-1H-indazol-5-yl)-5-fluoro-4-methylbenzamide



Example 137 was prepared by General Method D using using methyl (5-{5-
[(cyclopropylamino)carbonyl]-3-fluoro-2-methylphenyl}-1H-indazol-1-yl)acetate
5 (Example 123) and 1-cyclopropylmethanamine to give the title compound (0.008g).

LCMS: Rt 2.96min, MH+ 421.


Example 138: N-Cyclopropyl-3-fluoro-4-methyl-5-{1-[2-(4-morpholinyl)-2-oxoethyl]-1H-indazol-5-yl}benzamide



10 Example 138 was prepared by General Method D using using methyl (5-{5-
[(cyclopropylamino)carbonyl]-3-fluoro-2-methylphenyl}-1H-indazol-1-yl)acetate
(Example 123) and morpholine to give the title compound (0.001g).

LCMS: Rt 2.76min, MH+ 437.

15 **Example 139: N-Cyclopropyl-4-methyl-3-(1-{2-[(1-methylethyl)amino]-2-oxoethyl}-1H-indazol-5-yl)benzamide**




Example 139 was prepared by General Method D using methyl (5-{5-[(cyclopropylamino)carbonyl]-2-methylphenyl}-1H-indazol-1-yl)acetate (Example 117) and 2-propylamine to give the title compound (0.002g).

5

LCMS: Rt 2.82min, MH+ 391.

Example 140: N-Cyclopropyl-3-(1-{2-[(2,2-dimethylpropyl)amino]-2-oxoethyl}-1H-indazol-5-yl)-5-fluoro-4-methylbenzamide




10

Example 140 was prepared by General Method D using using methyl (5-{5-[(cyclopropylamino)carbonyl]-3-fluoro-2-methylphenyl}-1H-indazol-1-yl)acetate (Example 123) and 2,2-dimethyl-1-propanamine to give the title compound (0.008g).

LCMS: Rt 3.21min, MH+ 437.

15

Example 141: N-Cyclopropyl-3-{2-[(cyclopropylcarbonyl)amino]-1,3-benzothiazol-6-yl}-4-methylbenzamide



Example 141 was made by General Method A using *N*-(6-bromo-1,3-benzothiazol-2-yl)cyclopropanecarboxamide and *N*-cyclopropyl-4-methyl-3-(4,4,5,5-tetramethyl-1,3,2-dioxaborolan-2-yl)benzamide (Intermediate 5) to give the title compound
 5 (0.4mg).

LCMS: Rt 2.67min, MH⁺ 392.

Abbreviations

| | | |
|----|---------------------|---|
| 10 | Ac ₂ O | Acetic anhydride |
| | ^t BuONO | <i>t</i> -Butyl nitrite |
| | CDI | Carbonyldiimidazole |
| | DBU | 1,8-Diazabicyclo[5.4.0]undec-7-ene |
| | DIPEA | <i>N,N</i> -Diisopropylethylamine |
| 15 | DME | 1,2-Dimethoxyethane |
| | DMF | Dimethylformamide |
| | DMSO | Dimethylsulphoxide |
| | Et ₂ O | Ether |
| | EtOH | Ethanol |
| 20 | Hal | Halogen |
| | HATU | <i>O</i> -(7-Azabenzotriazol-1-yl)- <i>N,N,N',N'</i> -tetramethyluronium hexafluorophosphate |
| | HOBT | 1-Hydroxybenzotriazole |
| | KOAc | Potassium Acetate |
| 25 | MeCN | Acetonitrile |
| | MeOH | Methanol |
| | Ms | Mesyl |
| | NaO ^t Bu | Sodium <i>tert</i> -butoxide |
| | NaOEt | Sodium Ethoxide |
| 30 | NCS | <i>N</i> -Chlorosuccinimide |

| | |
|------------------------|---|
| PdCl ₂ dppf | [1,1'-bis(diphenylphosphino)ferrocene]dichloropalladium (II) complex with dichloromethane (1:1) |
| Rt | Retention Time |
| SPE | Solid phase extraction |
| 5 THF | Tetrahydrofuran |

BIOLOGICAL EXAMPLES

10 The activity of compounds of formula (I) as p38 inhibitors may be determined by the following *in vitro* assays:

Fluorescence anisotropy kinase binding assay

15 The kinase enzyme, fluorescent ligand and a variable concentration of test compound are incubated together to reach thermodynamic equilibrium under conditions such that in the absence of test compound the fluorescent ligand is significantly (>50%) enzyme bound and in the presence of a sufficient concentration (>10x K_i) of a potent inhibitor the anisotropy of the unbound fluorescent ligand is measurably different from the bound value.

20 The concentration of kinase enzyme should preferably be $\geq 1 \times K_f$. The concentration of fluorescent ligand required will depend on the instrumentation used, and the fluorescent and physicochemical properties. The concentration used must be lower than the concentration of kinase enzyme, and preferably less than half the kinase enzyme concentration. A typical protocol is:

25 All components dissolved in Buffer of final composition 62.5 mM HEPES, pH 7.5, 1.25 mM CHAPS, 1.25 mM DTT, 12.5 mM MgCl₂ 3.3% DMSO.

p38 Enzyme concentration: 12 nM

Fluorescent ligand concentration: 5 nM

Test compound concentration: 0.1 nM - 100 μ M


30 Components incubated in 30 μ l final volume in NUNC 384 well black microtitre plate until equilibrium reached (5-30 mins)

Fluorescence anisotropy read in LJI Acquest.

Definitions: K_i = dissociation constant for inhibitor binding

K_f = dissociation constant for fluorescent ligand binding

The fluorescent ligand is the following compound:



35

which is derived from 5-[2-(4-aminomethylphenyl)-5-pyridin-4-yl-1H-imidazol-4-yl]-2-chlorophenol and rhodamine green.

Results

The compounds described in the Examples were tested as described above and had IC₅₀ values of <10 μ M.